














Prenatal and postnatal exposures to endocrine disrupting chemicals and timing of pubertal onset in girls and boys: a systematic review and meta-analysis

C.S. Uldbjerg ^{1,2,†}, T. Koch ^{1,2,†}, Y.-H. Lim ^{3,4}, L.S. Gregersen ^{1,2},
C.S. Olesen ^{1,2}, A.-M. Andersson ^{1,2}, H. Frederiksen ^{1,2},
B.A. Coull ⁵, R. Hauser ⁶, A. Juul ^{1,2,7,*‡}, and E.V. Bräuner ^{1,2,*‡}

¹Department of Growth and Reproduction, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark ²The International Research and Research Training Centre in Endocrine Disruption of Male Reproduction and Child Health (EDMaRC), Rigshospitalet, University of Copenhagen, Copenhagen, Denmark ³Section of Environmental Health, Department of Public Health, University of Copenhagen, Copenhagen, Denmark ⁴Department of Preventive Medicine, Seoul National University College of Medicine, Seoul, Republic of Korea ⁵Department of Biostatistics, T.H. Chan School of Public Health, Harvard University, Cambridge, MA, USA ⁶Department of Environmental Health, T.H. Chan School of Public Health, Harvard University, Cambridge, MA, USA ⁷Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

*Correspondence address. Department of Growth and Reproduction, Rigshospitalet, Blegdamsvej 9, DK-2100 Copenhagen, Denmark.
Tel: +45-3545-5085; E-mail: anders.juul@regionh.dk  <https://orcid.org/0000-0002-0534-4350> (A.J.); Tel: +45-4242-8550;
E-mail: elvira.vaclavik.braeuner@regionh.dk  <https://orcid.org/0000-0001-9183-4796> (E.V.B.)

Submitted on August 26, 2021; resubmitted on January 25, 2022; editorial decision on February 07, 2022

TABLE OF CONTENTS

- Introduction
- Methods
 - Study design
 - Protocol and registration
 - Search strategy
 - Eligibility criteria
 - Exclusion criteria
 - Selection of literature
 - Assessment of quality
 - Data extraction and qualitative trend synthesis
 - Meta-analysis
- Results
 - Study base
 - Qualitative trend synthesis (n = 52 publications)
 - Meta-analysis (n = 23 publications)
- Discussion
 - Overall findings of the review
 - Strengths and limitations of the review
 - Conclusions and wider implications

[†]These authors contributed equally to this work.

[‡]These senior authors contributed equally to this work.

BACKGROUND: Globally, the ages at pubertal onset for girls and boys have been decreasing during recent decades, partly attributed to excess body fat accumulation. However, a growing body of literature has recognized that endocrine disrupting chemicals (EDCs) may play an important role in this global trend, but the association has not yet been fully established.

OBJECTIVE AND RATIONALE: EDCs can interfere with normal hormone function and metabolism and play a role in pubertal onset. We aimed to systematically identify and evaluate the current evidence on the timing of pubertal onset in girls and boys following prenatal or postnatal exposures to xenobiotic EDCs.

SEARCH METHODS: Following PRISMA guidelines, we performed a systematic literature search of original peer-reviewed publications in the PubMed database through a block search approach using a combination of index MeSH and free text search terms. Publications were considered if they covered biomarkers of prenatal or postnatal exposures to xenobiotic EDCs (European Commission's list of category I EDCs) measured in maternal or child biospecimen and pubertal onset defined by the progression of the following milestones (and assessed in terms of the following measures): menarche (age), thelarche (Tanner staging) and pubarche (Tanner staging), in girls, and genital stage (Tanner staging), testicular volume (ml) and pubarche (Tanner staging), in boys.

OUTCOMES: The literature search resulted in 703 references, of which we identified 52 publications fulfilling the eligibility criteria for the qualitative trend synthesis and 23 publications for the meta-analysis. The qualitative trend synthesis provided data on 103 combinations of associations between prenatal or postnatal exposure to EDC compounds groups and puberty outcomes and the meta-analysis enabled 18 summary risk estimates of meta-associations.

WIDER IMPLICATIONS: Statistically significant associations in the qualitative trend synthesis suggested that postnatal exposure to phthalates may be associated with earlier thelarche and later pubarche. However, we did not find consistent evidence in the meta-analysis for associations between timing of pubertal onset in girls and boys and exposures to any of the studied xenobiotic EDCs. We were not able to identify specific pre- or postnatal windows of exposure as particularly critical and susceptible for effects of EDCs. Current evidence is subject to several methodological challenges and inconsistencies and evidence on specific exposure-outcome associations remains too scarce to firmly confirm EDC exposure as a risk factor for changes in age of pubertal onset in the general child population. To create a more uniform foundation for future comparison of evidence and to strengthen pooled studies, we recommend the use of more standardized approaches in the choice of statistical analyses, with exposure transformations, and in the definitions and assessments of puberty outcomes. The impact of mixtures of EDC exposures on the association also remains unestablished and would be valuable to elucidate for prenatal and postnatal windows of exposure. Future large, longitudinal epidemiological studies are needed to clarify the overall association.

Key words: endocrine disrupting chemicals / puberty / menarche / thelarche / pubarche / genital stage / testicular volume / prenatal exposure / postnatal exposure

Introduction

Puberty marks the dynamic transition from childhood to adulthood, during which the child's body undergoes marked physical changes leading to reproductive maturity. In girls, pubertal development commonly starts with breast budding (thelarche) and the first appearance of pubic hair (pubarche) followed by the first menstrual bleeding (menarche) roughly 3 years later, while in boys, testicular enlargement (testicular volume) is usually the first sign of puberty followed by a gradual development of external genitals (genital stage) (Parent et al., 2003). But the onset of pubertal events, and the speed by which puberty progresses, is subject to large interindividual differences. The age of pubertal onset also varies significantly between girls and boys; girls tend to enter puberty at an earlier age (8–13 years) than boys (9–14 years) (Parent et al., 2003). Globally, the age of pubertal onset has, due to largely unknown reasons, decreased substantially during recent decades, especially among girls (Herman-Giddens et al., 1997, 2001; Kaplowitz and Oberfield, 1999; Sun et al., 2002; Wu et al., 2002; Aksglaede et al., 2009; Sørensen et al., 2010). This trend toward earlier puberty is of considerable concern, because earlier onset of pubertal indicators such as age at menarche in girls and voice break in boys have been linked to short- and long-term adverse physical and mental health outcomes including increased risk of obesity (Prentice and Viner, 2013), Type 2 diabetes (Elks et al., 2013; Day et al., 2015), breast cancer (Day et al., 2017) and prostate cancer (Bonilla et al.,

2016), as well as depression (Hamlat et al., 2014) and anxiety (Kaltiala-Heino et al., 2003).

The timing of pubertal onset is determined by a variety of genetic factors and environmental influences in addition to gender (Parent et al., 2003). While genetic factors are predominant in determining individual variations in pubertal timing, accounting for ~60% of variations (Sørensen et al., 2013), they cannot explain the observed trend and it is increasingly recognized that non-genetic life style factors like body fat as well as prenatal and postnatal exposures to endocrine disrupting chemicals (EDCs) may play a role in perturbations of pubertal onset (Lucaccioni et al., 2020). An EDC is defined as 'an exogenous substance or mixture that alters the functions of the endocrine system and consequently causes adverse health effects in an intact organism or its progeny' (Bundesumwelt and Kem, 1996). In general, EDCs can be divided into persistent and non-persistent compounds. The persistent EDC compounds are characterized by a long biological half-life period (months to a few years) as they accumulate in body tissue, whereas non-persistent EDC compounds have a relatively short half-life period (hours to days) in the human body (Domínguez-Romero and Scheringer, 2019). It is hypothesized that EDCs can interfere with normal hormone function and metabolism, which can either delay or accelerate the onset of puberty (Lucaccioni et al., 2020).

Several epidemiological studies and a recent systematic review have addressed this issue over the past 10 years. However, evidence of an association between EDCs and changes in pubertal development

remains largely inconclusive due to inconsistency in findings and reporting of both earlier and later pubertal onset (Lee *et al.*, 2019; Castiello and Freire, 2021). To date, no meta-analysis has been conducted and a systematic quantitative and qualitative evaluation of the current epidemiological evidence on effects of environmental endocrine disrupting xenobiotics has become timely. In this systematic review, we aimed to identify and evaluate current evidence on the timing of pubertal onset in girls and boys following prenatal or postnatal exposures to xenobiotic EDCs.

Methods

Study design

A systematic literature review and meta-analysis was performed in accordance with the PRISMA statement for reporting systematic reviews and meta-analyses of observational studies.

Protocol and registration

A review protocol was registered at PROSPERO.org (registration number CRD42021223868) prior to the study. The protocol was approved by all authors.

Search strategy

We performed a systematic search of peer-reviewed original literature in English published prior to 1 February 2021 to identify the available evidence on xenobiotic EDC exposures and changes in pubertal onset in girls and boys. The systematic literature search was conducted in the PubMed database through a block search approach using a combination of index MESH and free text search terms. The identified search terms were divided into two search blocks: the first block covering the exposure (xenobiotic EDCs) and the second block covering the outcome (timing of pubertal onset). We subsequently performed a hand search of the reference lists of the included publications. The search specifications and respective hits in each search block are provided in the search protocol (Supplementary Table S1).

Eligibility criteria

Publications were considered eligible if the following criteria were met.

Xenobiotic EDCs

Xenobiotic EDCs were defined as an exogenous man-made chemical or a mixture of chemicals that have been included in the European Commission's list of Category I EDCs (McCarthy, 2011) and measured in maternal (prenatal or proxy of prenatal) or child (postnatal) biospecimens. Throughout the manuscript, we use the term 'exposure' to represent the use of exposure biomarkers. The following 11 compound groups of EDCs were investigated in the included publications: (i) benzophenones and other UV filters (hereafter UV filters); (ii) bisphenol A (hereafter BPA); (iii) dioxins; (iv) flame retardants; (v) parabens; (vi) phthalates diesters and their metabolites (hereafter phthalates); (vii) polycyclic aromatic hydrocarbons (hereafter PAHs); (viii) polychlorinated biphenyl (hereafter PCBs); (ix) polychlorinated phenols/organochloride pesticides (hereafter polychlorinated phenols/

pesticides); (x) polyfluoroalkyl chemicals/per- and polyfluoroalkyl substances (hereafter PFCs/PFAS) and (xi) triclosan.

Pubertal onset

Pubertal onset was defined by the timing of the progression of the following pubertal milestones: menarche (age of first bleed), thelarche (Tanner staging), pubarche (Tanner staging), in girls, and genital stage (Tanner staging), testicular volume (ml) and pubarche (Tanner staging), in boys. Puberty outcomes using Tanner staging were determined by clinical assessments (visual inspection and/or palpitation) or self-report according to the Tanner scale, in which Stage 1 represents the prepubertal state and entrance into Stage 2 represents the onset of puberty (Marshall and Tanner, 1969). In girls, pubertal onset (B2+), a phenomenon also termed thelarche, is followed by subsequent breast enlargement corresponding to Tanner breast Stages B2 to B5. Menarche was assessed by retrospective self-report at the time of interview or via a diary completed prior to first bleed (exact age). In boys, pubertal onset is denoted by entrance into Tanner genital Stage 2 (G2+), followed by testicular/penis enlargement throughout genital Stage G2 to G5. Testicular volume was assessed by clinical assessment using orchidometer or ultrasound. For both sexes, appearance of pubic hair (curly pigmented hair) denotes Tanner pubic hair stage PH2 (a phenomenon termed pubarche), which usually occurs concurrently with gonadarche, i.e. breast or genital development. However, isolated pubic hair development may represent adrenal androgen secretion (a phenomenon termed adrenarche), which is not always associated with simultaneous activation of the hypothalamo-pituitary-gonadal hormone axis. However, it is not possible to distinguish gonadarche from adrenarche without thorough biochemical evaluation.

Exclusion criteria


Publications prior to 1 January 1990 or publications covering non-human studies, case-reports, experimental studies, reviews, conference proceedings/abstracts, letters, editorials and comments were excluded. We also excluded publications addressing natural estrogens (e.g. phytoestrogens or prescription hormones), publications not reporting risk estimates or duplicates repeating estimates reported in previous publications.

Selection of literature

We obtained 703 search references after removal of duplicates, including 11 references identified through hand searches and reference lists (Fig. 1). Titles and abstracts were screened by two authors (T.K. and C.S.O.) independently to assess eligibility. A total of 107 publications were considered eligible for full-text assessments. Among these, 55 publications were eventually excluded as they did not fulfill the defined eligibility criteria, such as exposure not measured in biospecimen ($n=9$) or outcome defined as idiopathic central precocious puberty ($n=16$). A total of 52 publications were included in the final review and qualitative trend synthesis. Of these, 23 publications were found eligible for the meta-analysis according to eligibility criteria.

Assessment of quality

The quality of the included publications was assessed to enable a critical interpretation of the respective study findings. All 52 publications



were evaluated for completeness of reporting and potential sources of bias using a standardized form adapted from [Bonzini et al. \(2007\)](#) and [Shamliyan et al. \(2010\)](#). (The evaluation form is available in [Supplementary Table SII](#).) The publications were evaluated independently by two authors (T.K. and C.S.O.) and any discrepancies in scoring were resolved through discussions with a third author (E.V.B.).

Completeness of reporting was assessed in the following 11 areas: (i) study design; (ii) sampling frame and procedures; (iii) inclusion and exclusion; (iv) population characteristics of exposed/unexposed or cases/referents; (v) response rate reported or implicitly given; (vi) methods for exposure measurement; (vii) method for outcome ascertainment; (viii) external quality assurance program of biochemical analyses; (ix) detection level and precision for biological samples; (x) statistical analysis and (xi) exposure-response. The 11 areas were equally weighted with the value one given for adequate reporting. We considered a sum of ≥ 8 as sufficient completeness of reporting.

Potential sources of bias were evaluated in seven areas, of which the following four areas were considered the most critical sources of bias and included for final assessment: (i) reporting of tested hypotheses; (ii) selection bias from loss of follow-up or lack of representativeness in a population sample; (iii) information bias related to outcome ascertainment and (iv) accounting for confounding. Each of the four areas were either rated as high risk, uncertain risk or low risk. Publications were considered at higher risk of bias if two or more of the given areas were identified as high risk of bias.

Data extraction and qualitative trend synthesis

Descriptive characteristics were extracted from each publication including study period, location, age at interview/examination, study design, sample size, biospecimens, outcome specifications, EDCs

examined, exposure window, etc. In a qualitative trend synthesis, the identified characteristics were presented in a table separately for girls and boys according to prenatal or postnatal exposure compounds along with a direction of the association between exposure and pubertal onset (regardless of statistical significance of the estimate) illustrated by arrows: downward pointing arrow indicating earlier onset; upward pointing arrow indicating later; and a horizontal arrow indicating no association (Tables II and III). The assigned assessment scores for completeness of reporting and risk of bias for individual publications were also presented. In a supplementary synthesis, the direction of all trends was presented according to each individual compound (rather than by publication) for girls and boys combined (Supplementary Table SIII). A risk of bias graph of all publications was also visualized in a separate figure (Fig. 2).

Meta-analysis

The meta-analysis provided summary risk estimates between specific exposures to EDCs and changes in pubertal onset in girls and boys. We applied a stepwise approach addressing three different strategies for grouping exposure compounds (Table I) separately for prenatal and postnatal exposures: (i) the associations according to compound groups of EDCs and timing of pubertal onset in girls and boys (main analysis); (ii) the associations according to persistent and non-persistent EDC groups and timing of pubertal onset in girls and boys; and (iii) the associations according to chemicals obtained on the EU REACH 2020 (European Chemicals Agency, 2021a) and/or 2021 Substances of Very High Concern (SVHC) lists (European Chemicals Agency, 2021b) and timing of pubertal onset in girls and boys.

The three strategies attempted to address the effects of groups and multiple combined EDC compounds that would be expected in real life, but we acknowledge that individual compounds within the classes may vary in potency. Further, the biological activity may vary, for instance with some EDCs being estrogenic, while others having been shown to possess anti-androgenic properties, or even both.

Publications were only included in the meta-analysis if the estimates were reported as a relative risk (RR; odds ratio, hazard ratio or risk

ratio) associated with exposure contrast (Hartemink *et al.*, 2006; Hagan *et al.*, 2011) and if the outcome was defined as Tanner stage ≥ 2 for thelarche (B2+), pubarche (PH2+) or genital stage (G2+); >3 ml for testicular volume; or yes/no for menarche.

Within each strategy, separate forest plots for each puberty outcome stratified by prenatal or postnatal exposure to specific compounds were created to illustrate summary RRs with 95% CIs. We standardized risk estimates to a common log scale ($\log_{10}(\text{RR})$) prior to analyses, as many EDC exposures had been reported as estimates based on various log scales (e.g. \log_2 , \log_{10} , natural \log_e) due to lack of normal distribution. We used a 10-fold exposure increase to identify small differences. Exposure-outcome summary risk estimates in the meta-analyses were only provided if multiple publications (at least two) were available and if three or more individual exposure-outcome estimates were available. We pooled the risk estimates by a random-effect meta-analysis and computed a common summary risk estimate by weighting the risk estimates with the inverse variance (within-study and between-studies) computed from the provided confidence limits. A formal measure of heterogeneity was provided using I^2 statistics. All statistical analyses were performed using R (version 3.6.0, R Foundation for Statistical Computing, Vienna, Austria) with a significance level of 0.05.

Results

Study base

We identified 52 publications (Blanck *et al.*, 2000; Vasiliu *et al.*, 2004; Warner *et al.*, 2004, 2020; Denham *et al.*, 2005; Ouyang *et al.*, 2005; Leijes *et al.*, 2008; Wolff *et al.*, 2008, 2010, 2014, 2015, 2017; Small *et al.*, 2009; Chen *et al.*, 2011; Christensen *et al.*, 2011; Den Hond *et al.*, 2011; Humblet *et al.*, 2011; Korrick *et al.*, 2011; Lopez-Espinosa *et al.*, 2011; Buttke *et al.*, 2012; Frederiksen *et al.*, 2012; Grandjean *et al.*, 2012; Mouritsen *et al.*, 2013; Ferguson *et al.*, 2014; Lam *et al.*, 2014; Watkins *et al.*, 2014, 2017a,b; Hou *et al.*, 2015; McGuinn *et al.*, 2015; Shi *et al.*, 2015; Su *et al.*, 2015; Windham *et al.*, 2015; Zhang




Table 1 List of specific chemical substances and abbreviations.

Individual EDC compounds	Abbreviation	EDC groupings	Included in meta-analysis		
			Strategy 1	Strategy 2	Strategy 3
			EDC grouping	Persistent/non-persistent	REACH 2020 and SVHC 2021
Benzophenone-2	BP-2	Benzophenones and other UV filters	×	× non-persistent	
Benzophenone-3	BP-3	Benzophenones and other UV filters	×	× non-persistent	
Ethyl-hexyl methoxycinnamate	EHMC	Benzophenones and other UV filters	×	× persistent	
4'-methoxyacetophenone	4'-MAP	Benzophenones and other UV filters	×	× persistent	
Ethylhexyl dimethyl PABA	OD-PABA	Benzophenones and other UV filters	×	× persistent	
Bisphenol A	BPA	Bisphenol A	×	× non-persistent	×
Bisphenol A-G	BPA-G	Bisphenol A	×	× non-persistent	×
Polychlorinated dibenzo-p-dioxins	PCDD	Dioxins			
Polychlorinated dibenzo-p-furans	PCDF	Dioxins			
2,3,7,8-Tetrachlorodibenzo-p-dioxins	TCDD	Dioxins		× persistent	×
Polybrominated biphenyl	PBB	Flame retardants			
2,4,4'-Tribromodiphenyl ether	BDE-28	Flame retardants	×	× persistent	×
2,2',4,4'-Tetrabromodiphenyl ether	BDE-47	Flame retardants	×	× persistent	×
2,2',4,4',5-Pentabromodiphenyl ether	BDE-99	Flame retardants	×	× persistent	×
2,2',4,4',6-Pentabromodiphenyl ether	BDE-100	Flame retardants	×	× persistent	×
2,2',4,4',5,5'-Hexabromodiphenyl ether	BDE-153	Flame retardants	×	× persistent	×
2,2',4,4',5,6'-Hexabromodiphenyl ether	BDE-154	Flame retardants	×	× persistent	×
Methyl paraben	MePB	Parabens		× non-persistent	
Propyl paraben	PrPB	Parabens		× non-persistent	×
Mono-methyl phthalate [DMP metabolite]	MMP	Phthalate diesters and their metabolites	×	× non-persistent	
Mono-ethyl phthalate [DEP metabolite]	MEP	Phthalate diesters and their metabolites	×	× non-persistent	
Mono-iso-butyl phthalate [DiBP metabolite]	MiBP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-(2-hydroxy-iso-butyl) phthalate [DiBP metabolite]	2OH-MiBP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-n-butyl phthalate [DnBP metabolite]	MnBP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-(3-hydroxybutyl) phthalate [DnBP metabolite]	3OH-MnBP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-benzyl phthalate [BBzP metabolite]	MBzP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-(2-ethyl-hexyl) phthalate [DEHP metabolite]	MEHP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-(2-ethyl-5-hydroxyhexyl) phthalate [DEHP metabolite]	5OH-MEHP	Alternative: MEHHP	×	× non-persistent	×
Mono-(2-ethyl-5-oxohexyl) phthalate [DEHP metabolite]	5oxo-MEHP	Alternative: MEOHP	×	× non-persistent	×
Mono-(2-ethyl-5-carboxypentyl) phthalate [DEHP metabolite]	5cx-MEPP	Alternative: MECPP	×	× non-persistent	×
Mono-3-carboxypropyl phthalate [DnOP metabolite, non-specific]	MCPP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-iso-nonyl phthalate [DiNP metabolite]	MiNP	Phthalate diesters and their metabolites	×	× non-persistent	×
Mono-hydroxy-iso-nonyl phthalate [DiNP metabolite]	OH-MiNP	Alternative: MHiNP	×	× non-persistent	×
Mono-oxo-iso-nonyl phthalate [DiNP metabolite]	oxo-MiNP	Alternative: MOiNP	×	× non-persistent	×

Continued

Table 1 Continued

Individual EDC compounds	Abbreviation	EDC groupings	Included in meta-analysis		
			Strategy 1	Strategy 2	Strategy 3
			EDC grouping	Persistent/non-persistent	REACH 2020 and SVHC 2021
Mono-carboxy-iso-octyl phthalate [DiNP metabolite]	cx-MiOP	Alternative: MCIOP	Phthalate diesters and their metabolites	× non-persistent	×
Mono-(hydroxy-iso-decyl) phthalate [DiDP metabolite]	OH-MiDP	Alternative: MHiDP	Phthalate diesters and their metabolites	× non-persistent	×
Mono-(oxo-iso-decyl) phthalate [DiDP metabolite]	oxo-MiDP	Alternative: MOiDP	Phthalate diesters and their metabolites	× non-persistent	×
Mono-(carboxy-iso-nonyl) phthalate [DiDP metabolite]	cx-MiNP	Alternative: MCiNP	Phthalate diesters and their metabolites	× non-persistent	×
Polychlorinated biphenyl	PCB	Polychlorinated biphenyl		× persistent	×
2,4-dichlorophenol	2,4-DCP	Polychlorinated phenols/organochloride pesticides		× non-persistent	
2,5-dichlorophenol	2,5-DCP	Polychlorinated phenols/organochloride pesticides		× non-persistent	
1,1-Dichloro-2,2-bis(p-chlorophenyl)ethylene	DDE	Polychlorinated phenols/organochloride pesticides		× persistent	×
2,2-Bis(4-chlorophenyl)-1,1-dichloroethylene	p,p'-DDE	Polychlorinated phenols/organochloride pesticides			
1,1'-(2,2,2-Trichloroethane-1,1-diyl)bis(4-chlorobenzene)	DDT	Polychlorinated phenols/organochloride pesticides		× persistent	×
1-chloro-4-[2,2,2-trichloro-1-(4-chlorophenyl)ethyl]benzene	p,p'-DDT	Polychlorinated phenols/organochloride pesticides		× persistent	×
Hexachlorobenzene	HCB	Polychlorinated phenols/organochloride pesticides		× persistent	×
Beta-hexachlorobenzene	β-HCB	Polychlorinated phenols/organochloride pesticides			
Beta-hexachlorocyclohexane	β-HCH	Polychlorinated phenols/organochloride pesticides			
3-phenoxybenzoic acid [metabolite of pyrethroid]	3-PBA	Polychlorinated phenols/organochloride pesticides			
Perfluorohexane sulfonate	PFHxS	PFCs or PFAS		× persistent	×
Perfluorononanoate	PFNA	PFCs or PFAS		× persistent	×
Perfluorooctanoic acid	PFOA	PFCs or PFAS		× persistent	×
Perfluorooctane sulfonic acid	PFOS	PFCs or PFAS		× persistent	×
Perfluorooctane sulfonamide	PFOSA	PFCs or PFAS		× persistent	×
2-(N-ethyl-perfluorooctane sulfonamido) acetate	Et-PFOSA-AcOH	PFCs or PFAS		× persistent	×
2-(N-methyl-perfluorooctane sulfonamido) acetate	Me-PFOSA-AcOH	PFCs or PFAS		× persistent	×
Triclosan	TCS	Triclosan	×	× non-persistent	

EDC, endocrine disrupting chemical.

Table II Study characteristics and qualitative trend synthesis for associations between prenatal or postnatal exposures to endocrine disrupting chemicals and menarche, M (age, n = 31 publications), thelarche, T (Tanner staging, n = 25 publications) and pubarche, P (Tanner staging, n = 23 publications) in girls.

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias ^S	Main meta-analysis
										↓ Earlier ↑ Later ↔ None				
Prenatal exposure														
Benzophenones Bisphenol A	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	172	Urine	BP-3	↓	↓	↑	10	0	No
	Berger et al. (2018)	USA	1999–2000	9	Longitudinal cohort	537	Urine	BPA	↑	↑*	↑	8	0	No
	Watkins et al. (2014)	Mexico	1997–2004	8–13	Longitudinal cohort	116	Urine	BPA	↑	↑	↓	7	1	No
	Watkins et al. (2017b)	Mexico	1997–2004	8–13	Longitudinal cohort	120	Urine	BPA	↑	↑	↓	8	1	No
Dioxins	Leijds et al. (2008) [#]	Netherlands	2005–2006	14–19	Longitudinal cohort	30	Breast milk	PCDF	↑*	↑*		8	1	No
	Warner et al. (2020)	Italy	1996	8–39	Longitudinal cohort	316	Serum	TCDD	↑			8	1	No
	Blanck et al. (2000) [#]	USA	1976–1979	5–24	Longitudinal cohort	327	Serum	PBB	↓	↓	↑*	10	1	No
	Harley et al. (2017)	USA	1999–2000	9	Longitudinal cohort	140	Serum	BDE-47	↑*	↑	↑	10	0	No
Flame retardants (Polybrominated diphenyl ethers, PBDE)								BDE-99	↑*	↔	↔			
								BDE-100	↑*	↑	↔			
								BDE-153	↑*	↑	↑			
								ΣPBDE	↑*	↑	↑			
Parabens	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	172	Urine	MePB	↓	↓	↑	10	0	No
								PrPB	↑	↑	↑			
	Berger et al. (2018)	USA	1999–2000	9	Longitudinal cohort	537	Urine	MBzP	↑	↑*	↑	8	0	No
								MCP	↑	↑	↑			
Phthalate diesters and their metabolites								cx-MiOP	↑	↑	↑			
								cx-MINP	↑	↑	↑			
								ΣDEHPm**	↑*	↑*	↑			
	Cathey et al. (2020)	Mexico	1997–2004	8–14	Longitudinal cohort	103	Urine	MEP	↓	↓	↓	8	1	No
								MIBP	↔	↔	↑			
								MnBP	↓	↓	↓			
								MBzP	↓	↓*	↓			
								MEHP	↓	↓	↓			
								5OH-MEHP	↓	↓*	↓			
								5oxo-MEHP	↓	↓*	↓			
								5cx-MEPP	↓	↓*	↓			
								MCP	↓	↓	↓			
								ΣDEHPm**	↓	↓*	↓			
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	172	Urine	MEP	↓	↓	↓*	10	0	No
								MIBP	↓	↓	↓			
								MnBP	↓	↓	↓			

Continued

Table II Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias ^S	Main meta-analysis
									↓ Earlier ↑ Later ↔ None					
Polychlorinated biphenyl (PCB)	Watkins et al. (2014)	Mexico	1997–2004	8–13	Longitudinal cohort	116	Urine	MEP	↑	↑	↑	7	1	Yes ^(M,T,P)
								MIBP	↑	↑	↑			
								MnBP	↑	↑	↑			
								MnBP	↑	↑	↑			
								MBzP	↑	↑	↑			
								MEHP	↑	↑	↑			
								5OH-MEHP	↑	↑	↑			
								5oxo-MEHP	↑	↑	↑			
								5ox-MEPP	↑	↑	↑			
								MCPP	↑	↑	↑			
								MEP	↑	↑	↑	8	1	Yes ^(M,T,P)
								MIBP	↑	↑	↑			
Polychlorinated phenols/organochloride pesticides	Watkins et al. (2017b)	Mexico	1997–2004	8–13	Longitudinal cohort	120	Urine	MIBP	↑	↑	↑			
								MnBP	↑	↑	↑			
								MnBP	↑	↑	↑			
								MBzP	↑	↑	↑			
								MEHP	↑	↑	↑			
								MCPP	↑	↑	↑			
								ΣDEHPm**	↑	↑	↑			
								ΣPCB	↑	↑	↑	10	1	No
								2,4-DCP	↑	↑	↑	10	0	No
								2,5-DCP	↑	↑	↑			
								p,p'-DDT	↑	↑	↑	10	1	No
								HCb	↑	↑	↑			
Polyfluoroalkyl chemicals/per- and polyfluoroalkyl substances	Namulanda et al. (2017)	England	1991–1992	8–17	Nested case-control	174/195	Urine	β-HCB	↑	↑	↑			
								D-atrazine	↑	↑	↑	10	1	No
								DDE	↑	↑	↑	10	1	No
								Carboxylates	↑	↑	↑	9	1	No
								PFHxS	↑	↑	↑			
								PFNA	↑	↑	↑			
								PFOA	↑	↑	↑			
								PFOS	↑	↑	↑			
								PFOSA	↑	↑	↑			
								Et-PFOSA-AcOH	↑	↑	↑			
								Me-PFOSA-AcOH	↑	↑	↑			
								sulfonamide esters	↑	↑	↑			
Triclosan	Ernst et al. (2019) ^{††}	Denmark	1996–2002	11	Longitudinal cohort	555	Serum	PFOA	↑	↑	↑	10	1	No
								PFOS	↑	↑	↑			
								TCS	↑	↑	↑	10	0	No
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	172	Urine							

Continued

Table II Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias ^S	Main meta-analysis
									↓ Earlier ↑ Later ↔ None					
Postnatal exposure														
Benzophenones and other UV filters	Binder <i>et al.</i> (2018)	Chile	2009	7–10	Longitudinal cohort	200	Urine	BP-3	↓			8	0	Yes ^(M)
	Buttke <i>et al.</i> (2012)	USA	2003–2008	12–16	Cross-sectional	461	Urine	BP-3	↑			9	0	Yes ^(M)
	Harley <i>et al.</i> (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	BP-3	↓	↓		10	0	No
	Huang <i>et al.</i> (2020) ^(baseline)	China	2011	7–15	Cross-sectional	244	Urine	BP-2	↓	↑	↔	8	0	Yes ^(M)
								BP-3	↓	↑	↓			
								EHMC	↓	↑	↔			
								4'-MAP	↓	↑	↓			
								OD-PABA	↑	↓	↓			
	Huang <i>et al.</i> (2020) ^(18 months follow-up)		+18 months	+18 months		176		BP-2	↓	↔	↓			
								BP-3	↔	↓	↑			
EHMC								↑	↔	↑				
4'-MAP								↑	↔	↓				
OD-PABA								↑	↓	↓	↑*			
Bisphenol A	Wolff <i>et al.</i> (2010)	USA	2004–2007	6–8	Longitudinal cohort	1151	Urine	BP-3	↓	↓	↓	8	0	No
	Wolff <i>et al.</i> (2015)	USA	2004–2007	6–8	Longitudinal cohort	1170	Urine	BP-3		↑*	↔	8	1	No
	Wolff <i>et al.</i> (2017)	USA	2004–2007	6–8	Longitudinal cohort	1051	Urine	BP-3	↑			8	1	Yes ^(M)
	Binder <i>et al.</i> (2018)	Chile	2009	7–10	Longitudinal cohort	200	Urine	BPA	↑			8	0	Yes ^(M)
	Buttke <i>et al.</i> (2012)	USA	2003–2008	12–16	Cross-sectional	461	Urine	BPA	↑			9	0	Yes ^(M)
	Howland <i>et al.</i> (2020) [#]	UK	2009–2011	5–16	Cross-sectional	348	Urine	BPA-G	↓	↓		8	1	Yes ^(T)
	Kasper-Sonnenberg <i>et al.</i> (2017) [#]	Germany	2009–2010	8–10	Longitudinal cohort	222	Urine	BPA	↓	↑	↓	7	1	Yes ^(M)
	McGuinn <i>et al.</i> (2015)	USA	2003–2010	12–19	Cross-sectional	987	Urine	BPA	↑	↑		9	0	Yes ^(M)
	Miao <i>et al.</i> (2017)	China	2011	9–18	Cross-sectional	655	Urine	BPA	↑*	↑	↓	8	0	No
	Watkins <i>et al.</i> (2014)	Mexico	1997–2004	8–13	Longitudinal cohort	116	Urine	BPA	↑	↑	↑	7	1	Yes ^(M,T,P)
Dioxins	Wolff <i>et al.</i> (2008)	USA	1996–1997	9	Cross-sectional	186	Urine	BPA	↑	↑	↑	8	0	Yes ^(T,P)
	Wolff <i>et al.</i> (2010)	USA	2004–2007	6–8	Longitudinal cohort	1151	Urine	BPA		↑	↔	8	0	No
	Wolff <i>et al.</i> (2015)	USA	2004–2007	6–8	Longitudinal cohort	1170	Urine	BPA		↑	↓	8	1	Yes ^(T,P)
	Wolff <i>et al.</i> (2017)	USA	2004–2007	6–8	Longitudinal cohort	1051	Urine	BPA	↓			8	1	Yes ^(M)
	Warner <i>et al.</i> (2004)	Italy	1996–1998	NR	Retrospective cohort	282	Serum	TCDD	↑			9	0	No
	Attfield <i>et al.</i> (2019)	USA	2004–2007	6–8	Longitudinal cohort	1257	Serum	Σ PBDE	↑			10	0	No
	Chen <i>et al.</i> (2011)	USA	2003–2004	12–19	Cross-sectional	271	Serum	BDE-28	↑*			9	0	Yes ^(M)
								BDE-47	↑*					
								BDE-99	↑*					
								BDE-100	↑*					
BDE-153								↑*						
BDE-154								↑*						
Σ PBDE								↑*						
Continued														

Continued

Table II Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias ^S	Main meta-analysis
									↑ Earlier ↓ Later ↔ None					
	Harley et al. (2017)	USA	1999–2000	9	Longitudinal cohort	266	Serum	BDE-47 BDE-99 BDE-100 BDE-153 Σ PBDE	↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑	↔ ↔ ↑ ↑ ↔	10	I	Yes ^(M)
	Windham et al. (2015)	USA	2004–2007	6–8	Longitudinal cohort	645	Serum	Σ PBDE	↑*	↑*	↑*	8	0	No
	Binder et al. (2018)	Chile	2009	7–10	Longitudinal cohort	200	Urine	MePB PrPB	↔ ↑	↔ ↑	↔ ↔	8	0	No
	Buttke et al. (2012)	USA	2003–2008	12–16	Cross-sectional	461	Urine	Σ parabens	↑	↑	↔	9	0	No
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	MePB PrPB	↓* ↑	↓* ↑	↓* ↑	10	0	No
	Wolff et al. (2010)	USA	2004–2007	6–8	Longitudinal cohort	1151	Urine	Σ parabens	↑	↑	↑	8	0	No
	Wolff et al. (2015)	USA	2004–2007	6–8	Longitudinal cohort	1170	Urine	Σ parabens	↑	↑	↑	8	I	No
	Wolff et al. (2017)	USA	2004–2007	6–8	Longitudinal cohort	1051	Urine	Σ parabens	↑	↑	↑	8	I	No
	Binder et al. (2018)	Chile	2009	7–10	Longitudinal cohort	200	Urine	MMP MEP MIBP 2OH-MIBP MnBP MBzP MEHP 5OH-MEHP 5oxo-MEHP 5ox-MEPP MCP α-MiOP α-MINP Σ DEHPm**	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔ ↔	8	0	Yes ^(M)
	Buttke et al. (2012)	USA	2003–2008	12–16	Cross-sectional	461	Urine	Σ phthalates	↑	↑	↑	9	0	No
	Frederiksen et al. (2012)	Denmark	2006–2008	6–19	Cross-sectional	725	Urine	MEP MIBP+MnBP MBzP Σ DEHPm** Σ DINPm** Σ phthalates Σ corr.phthalates	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	9	I	No

Continued

Table II Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias [§]	Main meta-analysis
										↓ Earlier ↑ Later ↔ None				
	Hou et al. (2015) [#]	Taiwan	2012–2003	7–15	Cross-sectional	270	Urine	MMP MEP MIBP MnBP MBzP MEHP 5OH-MEHP 5oxo-MEHP 5cx-MEPP ΣDEHPm** Σlow-MWP Σhigh-MWP Σphthalates	↑ ↔ ↔ ↓ ↔ ↑ ↓ ↓ ↓ ↑ ↑ ↑ ↑	↔ ↑ ↔ ↓ ↔ ↑ ↓ ↓ ↓ ↑ ↑ ↑ ↑	9	I	No	
	Kasper-Sonnenberg et al. (2017) [#]	Germany	2009–2010	8–10	Longitudinal cohort	222	Urine	MMP MEP MIBP 2OH-MiBP MnBP 3OH-MnBP MBzP MEHP 5OH-MEHP 5oxo-MEHP 5cx-MEPP MINP OH-MINP oxo-MINP cx-MiOP OH-MiDP oxo-MiDP	↑ ↑ ↓ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↓ ↓ ↓ ↓ ↑	↑ ↑* ↑ ↑ ↓ ↑ ↑ ↑ ↑ ↑ ↑ ↓ ↑ ↑ ↑ ↑ ↑	7	I	Yes ^(M)	
	Mouritsen et al. (2013)	Denmark	2006–2010	6–13	Longitudinal cohort	84	Urine	MEP MiBP + MnBP MBzP ΣDEHPm** ΣDINPm** Σcorr.phthalates	↑ ↑ ↑ ↓ ↑ ↑	↔ ↑ ↓ ↔ ↔ ↑	8	I	No	

Continued

Table III Continued

Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias [§]	Main meta-analysis						
								↓ Earlier ↑ Later											
								↔ None											
Shi et al. (2015)	China	2010	7–14	Cross-sectional	186	Urine	MMP	↑	↑	↑	10	1	Yes ^(M)						
							MEP	↑	↔	↑									
							MnBP	↑	↑	↑									
							MEHP	↑	↓	↑									
							5OH-MEHP	↑	↓	↑									
							5oxo-MEHP	↑	↓	↑									
							ΣDEHPm**	↑	↓	↑									
Watkins et al. (2014)	Mexico	1997–2004	8–13	Longitudinal cohort	116	Urine	MEP	↓	↓	↓	7	1	Yes ^(M,T,P)						
							MIBP	↓	↓	↓									
							MnBP	↓	↓	↓									
							MBzP	↓	↓	↓									
							MEHP	↑	↓	↓									
							5OH-MEHP	↓	↓	↓									
							5oxo-MEHP	↓	↓	↓									
							5cx-MEPP	↓	↓	↓									
Wolff et al. (2010)	USA	2004–2007	6–8	Longitudinal cohort	1151	Urine	Σlow-MWVP	↓	↓	↓	8	0	No						
							Σhigh-MWVP	↓	↓	↓									
Wolff et al. (2014)	USA	2004–2007	6–8	Longitudinal cohort	1170	Urine	MEP	↓	↓	↑	8	0	Yes ^(r,p)						
							MIBP	↑	↑	↑									
							MnBP	↑	↑	↑									
							MBzP	↑	↑	↑									
							MEHP	↑	↑	↑									
							5OH-MEHP	↑	↑	↑									
							5oxo-MEHP	↓	↓	↑									
							5cx-MEPP	↓	↓	↑									
							MCPP	↑	↑	↑									
							ΣDEHPm**	↓	↓	↑									
							Σlow-MWVP	↓	↓	↑									
							Σhigh-MWVP	↓	↓	↑									
							Wolff et al. (2017)	USA	2004–2007	6–8	Longitudinal cohort	1051	Urine	MEP	↓	↓	↑	8	1
MIBP	↑	↑	↑																
MnBP	↑	↑	↑																
MBzP	↑	↑	↑																
MEHP	↑	↑	↑																
5OH-MEHP	↑	↑	↑																
5oxo-MEHP	↑	↑	↑																
5cx-MEPP	↓	↓	↑																
MCPP	↑	↑	↑																
ΣDEHPm**	↑	↑	↑																
Σlow-MWVP	↑	↑	↑																
Σhigh-MWVP	↑	↑	↑																

Continued

Table II Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias [§]	Main meta-analysis
									↓ Earlier ↑ Later ↔ None					
Polyaromatic hydrocarbons (PAH)	Zhang et al. (2015) ^(baseline)	China	2011	7–14	Cross-sectional	251	Urine	MMP	↑	↑	↑	10	0	Yes ^(M,T,P)
								MEP	↑	↑	↑			
								MnBP	↑	↑	↑			
								MEHP	↑	↑	↑			
								5OH-MEHP	↑	↑	↑			
								5oxo-MEHP	↑	↑	↑			
								ΣDEHPm**	↑	↑	↑			
						208		MMP	↑	↑	↑			
								MEP	↑	↑	↑			
								MnBP	↑	↑	↑			
								MEHP	↑	↑	↑			
								5OH-MEHP	↑	↑	↑			
								5oxo-MEHP	↑	↑	↑			
								ΣDEHPm**	↑	↑	↑			
Polyaromatic hydrocarbons (PAH)	Dobrca et al. (2020)	USA	2005–2006	6–8	Longitudinal cohort	404	Urine	Σfluorene	↑	↑	↑	8	0	No
								1-hydroxypyrene	↑	↑	↑			
								2-naphthol	↑	↑	↑			
								ΣPAH	↑	↑	↑			
								Σphenanthrene	↑	↑	↑			
								ΣPCB	↑	↑	↑	10	0	No
								estrogenic ΣPCB	↑	↑	↑	7	0	No
Polychlorinated biphenyl (PCB)	Attfield et al. (2019)	USA	2004–2007	6–8	Longitudinal cohort	1257	Serum	ΣPCB	↑	↑	↑	9	1	No
	Denham et al. (2005)	Canada	1996–2000	10–17	Cross-sectional	138	Serum	ΣPCB	↑	↑	↑	8	0	No
	Den Hond et al. (2011)	Belgium	2003–2004	14–15	Cross-sectional	636	Serum	ΣPCB	↑	↑	↑	8	0	No
	Windham et al. (2015)	USA	2004–2007	6–8	Longitudinal cohort	645	Serum	ΣPCB	↑	↑	↑	8	0	No
	Wolff et al. (2008)	USA	1996–1997	9	Cross-sectional	186	Urine	ΣPCB	↑	↑	↑	10	0	No
	Attfield et al. (2019)	USA	2004–2007	6–8	Longitudinal cohort	1257	Serum	ΣOCP	↑	↑	↑	8	0	No
	Binder et al. (2018)	Chile	2009	7–10	Longitudinal cohort	200	Urine	2,4-DCP	↑	↑	↑	8	0	Yes ^(M)
Polychlorinated phenols/organochloride pesticides (OCP)	Burtke et al. (2012)	USA	2003–2008	12–16	Cross-sectional	461	Urine	2,5-DCP	↑	↑	↑	9	0	Yes ^(M)
								2,4-DCP	↑	↑	↑			
								2,5-DCP	↑	↑	↑			
								Σphenols	↑	↑	↑			
	Denham et al. (2005)	Canada	1996–2000	10–17	Cross-sectional	138	Serum	p,p'-DDE	↑	↑	↑	7	0	No
								HCB	↑	↑	↑			
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	2,4-DCP	↑	↑	↑	10	0	No
								2,5-DCP	↑	↑	↑			
	Hou et al. (2015) [#]	Taiwan	2012–2013	7–15	Cross-sectional	270	Urine	nonylphenol	↑	↑	↑	9	1	No
	Ouyang et al. (2005)	China	1996–1998	20–34	Cross-sectional	466	Serum	DDT	↑	↑	↑	10	1	No

Continued

Table II Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	M	T	P	CR	Bias [§]	Main meta-analysis
									↓ Earlier ↑ Later ↔ None					
	Windham et al. (2015)	USA	2004–2007	6–8	Longitudinal cohort	645	Serum	Σ OCP		↑	↑	8	0	No
	Wolff et al. (2008)	USA	1996–1997	9	Cross-sectional	186	Serum	DDE		↓	↓	8	0	No
	Wolff et al. (2010)	USA	2004–2007	6–8	Longitudinal cohort	1151	Urine	2,5-DCP		↑	↑	8	0	No
	Wolff et al. (2015)	USA	2004–2007	6–8	Longitudinal cohort	1170	Urine	2,5-DCP		↓*	↓*	8	1	No
	Wolff et al. (2017)	USA	2004–2007	6–8	Longitudinal cohort	1051	Urine	2,5-DCP	↓			8	1	Yes ^(M)
	Ye et al. (2017) [#]	China	2014–2015	9–15	Cross-sectional	305	Urine	3-PBA	↑*	↑	↑*	9	0	No
Polyfluoroalkyl chemicals/ per- and polyfluoroalkyl substances	Lopez-Espinosa et al. (2011)	USA	2005–2006	8–18	Cross-sectional	2931	Serum	PFOA	↑*			11	1	No
								PFOA	↑*					
								PFOA	↑*					
Triclosan	Binder et al. (2018)	Chile	2009	7–10	Longitudinal cohort	200	Urine	TCS	↔			8	0	Yes ^(M)
	Burtke et al. (2012)	USA	2003–2008	12–16	Cross-sectional	461	Urine	TCS	↑			9	0	Yes ^(M)
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	TCS	↑	↔	↑	10	0	No
	Wolff et al. (2010)	USA	2004–2007	6–8	Longitudinal cohort	1151	Urine	TCS		↓	↓	8	0	No
	Wolff et al. (2015)	USA	2004–2007	6–8	Longitudinal cohort	1170	Urine	TCS		↓*	↓*	8	1	No
	Wolff et al. (2017)	USA	2004–2007	6–8	Longitudinal cohort	1051	Urine	TCS	↓			8	1	Yes ^(M)
								TCS				8	1	

↑ Later onset; ↓ Earlier onset; ↔ none detected

[#] Self-reported thelarche or pubarche rather than clinical examination (age at menarche is self-reported in all publications)

[§] CR: Completeness of reporting ≥ 8 was considered sufficient; Bias and confounding: 1 = potential risk of bias, 0 = lower risk of bias

*P-value < 0.05

^(M,1,P) Included in meta-analysis for menarche (M); thelarche (T); pubarche (P)

** Sums of selected phthalate metabolites and substitutes:

Σ DEHPm: molar sum of MEHP, 5OH-MEHP, 5oxo-MEHP, 5cx-MEPP and 2cx-MMHP expressed as DEHP

Σ DINPm: molar sum of MINP, OH-MINP, 6oxo-MINP and 6cx-MIOP expressed as DINP

low/high MWP: low/high molecular weight phthalates

EDC, endocrine disrupting chemical.

Table III Study characteristics and qualitative trend synthesis for associations between prenatal or postnatal exposures to endocrine disrupting chemicals and genital stage, GS (Tanner staging, n = 17), testicular volume, TV (ml, n = 15) and pubarche, P (Tanner staging, n = 20) in boys.

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	GS	TV	P	CR	Bias ^S	Main meta-analysis
Prenatal exposure														
Benzophenones	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	BP-3	↑	↑	↑	10	0	No
	Berger et al. (2018)	USA	1999–2000	9	Longitudinal cohort	537	Urine	BPA	↓*	↓*	↓*	8	0	No
	Ferguson et al. (2014)	Mexico	1994–2004	8–14	Longitudinal cohort	107	Urine	BPA	↑	↑	↑	7	1	No
	Watkins et al. (2017a)	Mexico	1997–2004	8–14	Longitudinal cohort	109	Urine	BPA	↑	↑	↑	7	1	No
Dioxins	Humblet et al. (2011) ^S	Russia	2003–2005	8–9	Longitudinal cohort	424	Serum	Σ _{Toxic Equiv. Quotient}	↑	↑	↑	9	0	No
	Small et al. (2009) [#]	USA	1973–1974	5–30	Longitudinal cohort	464	Serum	PBB	↑*	↑*	↑*	10	1	No
	Harley et al. (2017)	USA	1999–2000	9	Longitudinal cohort	117	Serum	BDE-47	↔	↔	↔	10	0	No
								BDE-99	↑	↑	↑			
Flame retardants (Polybrominated diphenyl ethers, PBDE)	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	BDE-100	↔	↔	↔			
								BDE-153	↑	↑	↑			
								Σ PBDE	↔	↔	↔			
								MePB	↔	↔	↔	10	0	No
Parabens	Berger et al. (2018)	USA	1999–2000	9	Longitudinal cohort	537	Urine	PrPB	↑	↑	↑	8	0	No
								MBzP	↑*	↑*	↑*			
								MCPP	↑*	↑*	↑*			
								α-MiOP	↑*	↑*	↑*			
Phthalate diesters and their metabolites	Cathey et al. (2020)	Mexico	1997–2004	8–14	Longitudinal cohort	91	Urine	α-MiNP	↑*	↑*	↑*			
								Σ DEHPm**	↑	↑	↑			
								MEP	↑	↑	↑	8	1	No
								MIBP	↑	↑	↑			
								MnBP	↑	↑	↑			
								MBzP	↑	↑	↑			
								MEHP	↑	↑	↑			
								5OH-MEHP	↑	↑	↑			
								5oxo-MEHP	↑	↑	↑			
								5α-MEPP	↑	↑	↑			
								MCPP	↑	↑	↑			
								Σ DEHPm**	↑	↑	↑			

Continued

Table III Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	GS	TV	P	CR	Bias ^S	Main meta-analysis	
										↓ Earlier ↑ Later ↔ None					
Polychlorinated biphenyl (PCB)	Ferguson et al. (2014)	Mexico	1994–2004	8–14	Longitudinal cohort	107	Urine	MEP MIBP MnBP MBzP MEHP 5OH-MEHP 5oxo-MEHP 5ox-MEPP MCPp	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	7	1	Yes ^(GS,TV,P)	
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	MEP MIBP MnBP	↑ ↑ ↑	↑ ↑ ↑	↑ ↑ ↑	10	0	No	
	Su et al. (2015)	Taiwan	2000–2001	8	Longitudinal cohort	59	Urine	MMP MEP MnBP MBzP MEHP 5OH-MEHP 5oxo-MEHP ΣDEHPm**	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑ ↑ ↑ ↑	9	1	No	
	Watkins et al. (2017a)	Mexico	1997–2004	8–14	Longitudinal cohort	109	Urine	MEP MBzP MCPp MIBP+MnBP ΣDEHPm**	↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑	↑ ↑ ↑ ↑ ↑	7	1	Yes ^(GS,TV,P)	
	Grandjean et al. (2012)	Faroe Islands	1986–1987	14	Longitudinal cohort	438	Coord blood	ΣPCB	↑	↑	↑	9	0	No	
	Humblet et al. (2011)	Russia	2003–2005	8–9	Longitudinal cohort	424	Serum	ΣPCB	↑ [*]	↑ ↑	↑ ↑	9	0	No	
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	2,4-DCP 2,5-DCP	↑ ↑	↑ ↑	↑ ↑	10	0	No	
	Ernst et al. (2019) [#]	Denmark	1996–2002	11	Longitudinal cohort	565	Serum	PFOA PFOS	↑ ↑	↑ ↑	↑ ↑	10	1	No	
	Triclosan	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	155	Urine	TCS	↑	↑	↑	10	0	No

Continued

Table III Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	GS	TV	P	CR	Bias ^{\$}	Main meta-analysis
Postnatal exposure														
Benzophenones and other UV filters	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	139	Urine	BP-3	↓	↓	↓	10	0	No
	Huang et al. (2020) (baseline)	China	2011	7–15	Cross-sectional	277	Urine	BP-2	↑	↑	↑	8	0	No
								BP-3	↑	↑	↑			
								EHMC	↑*	↑*	↑			
								4'-MAP	↑*	↑	↑			
Bisphenol A	Huang et al. (2020) (18 months follow-up)		+18 months	+18 months		151		OD-PABA	↑	↑	↑			
								BP-2	↓	↓	↓			
								BP-3	↑	↑*	↑			
								EHMC	↑*	↑*	↓			
								4'-MAP	↓	↑	↓			
	Ferguson et al. (2014)	Mexico	1994–2004	8–14	Longitudinal cohort	113	Urine	OD-PABA	↓	↑	↓	7	1	No
								BPA	↑	↓	↓	7	1	No
								BPA		↑	↑	10	0	No
								BPA	↓	↓	↑	9	0	No
								∑ Toxic Equiv. Quotient	↑*	↑*	↑			
Dioxins	Burns et al. (2016)	Russia	2003–2005	8–9	Longitudinal cohort	473	Serum	∑ Dioxin Like Compounds	↓	↑*	↓			
								TCDD	↓	↑*	↓	9	0	No
Flame retardants (Polybrominated diphenyl ethers, PBDE)	Harley et al. (2017)	USA	1999–2000	9	Longitudinal cohort	266	Serum	PCDD	↓	↑	↑			
								PCDF	↑	↑	↑	10	0	No
								BDE-47	↔	↔	↔			
								BDE-99	↔	↔	↔			
								BDE-100	↓	↓	↓			
								BDE-153	↓	↓	↓			
								∑ PBDE	↓	↔	↔			
Parabens	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	139	Urine	MePB	↓	↑	↑	10	0	No
								PrPB	↓*	↓	↓			
Phthalate diesters and their metabolites	Ferguson et al. (2014)	Mexico	1994–2004	8–14	Longitudinal cohort	113	Urine	MEP	↓	↓	↓	7	1	Yes ^(14, 16)
								MIBP	↑	↑	↑			
								MnBP	↓	↓*	↓			
								MBzP	↓	↑	↑			
								MEHP	↓	↓	↓			
								5OH-MEHP	↓	↑	↑			
								5oxo-MEHP	↓	↑	↑			
								5cx-MEPP	↑	↑	↑			
								MCPP	↓	↑	↓			
									↑	↑	↑			

Continued

Table III Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	GS			CR	Bias ^{\$}	Main meta-analysis
									↓ Earlier	↑ Later	↔ None			
	Hou et al. (2015) [#]	Taiwan	2012–2013	7–15	Cross-sectional	270	Urine	MMP MEP MIBP MnBP MBzP MEHP 5OH-MEHP 5oxo-MEHP 5cx-MEPP ΣDEHPm** Σlow-MWP Σhigh-MWP Σphthalates	↑*	↑	↔	9	I	No
	Kasper-Sonnenberg et al. (2017) [#]	Germany	2009–2010	8–10	Longitudinal cohort	250	Urine	MMP MEP MIBP 2OH-MIBP MnBP 3OH-MnBP MBzP MEHP 5OH-MEHP 5oxo-MEHP 5cx-MEPP MINP OH-MINP oxo-MINP cx-MIOP OH-MIDP oxo-MIDP	↑	↑	↑	7	I	No
	Mouritsen et al. (2013)	Denmark	2006–2010	6–13	Longitudinal cohort	84	Urine	MEP MIBP+MnBP MBzP ΣDEHPm** ΣDINPm** Σcorr.phthalates	↑	↑	↑	8	I	No

Continued

Table III Continued

	Author, year	Location	Study period	Age at interview (years)	Study design	Sample size	Matrix	EDC	GS	TV	P	CR	Bias [§]	Main meta-analysis
									↓ Earlier ↑ Later ↔ None					
	Shi et al. (2015)	China	2010	7–14	Cross-sectional	172	Urine	MMP MEP MnBP MEHP 5OH-MEHP 5oxo-MEHP ΣDEHPm**		↑ ↓ ↑* ↓ ↑ ↑ ↑	↑ ↓ ↑ ↓ ↑ ↑ ↑	I 0		No
	Zhang et al. (2015) (baseline)	China	2011	7–14	Cross-sectional	252	Urine	MMP MEP MnBP MEHP 5OH-MEHP 5oxo-MEHP ΣDEHPm**		↑ ↓ ↑ ↓ ↑ ↑ ↑	↑ ↓ ↑ ↓ ↑ ↑ ↑	I 0		Yes ^(TV, P)
	Zhang et al. (2015) (18 months follow-up)		+18 months	+18 months		222		MMP MEP MnBP MEHP 5OH-MEHP 5oxo-MEHP ΣDEHPm**		↑ ↓ ↑ ↓ ↑ ↑ ↑	↑ ↓ ↑ ↓ ↑ ↑ ↑			
Polychlorinated biphenyl (PCB)	Burns et al. (2016)	Russia	2003–2005	8–9	Longitudinal cohort	473	Serum	Σnon-dioxin-like PCB	↓	↑	↑	9	0	No
	Den Hond et al. (2011)	Belgium	2003–2004	14–15	Cross-sectional	767	Serum	ΣPCB	↑	↑	↑	9	I	No
	Korrick et al. (2011)	Russia	2003–2005	8–9	Longitudinal cohort	453	Serum	ΣPCB	↓	↑	↑	9	0	No
Polychlorinated phenols/ organochloride pesticides	Den Hond et al. (2011)	Belgium	2003–2004	14–15	Cross-sectional	767	Serum	p,p'-DDE HCB	↓ ↑	↑ ↑	↑ ↑	9 I		No
	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	139	Urine	2,4-DCP 2,5-DCP	↓ ↑	↑ ↑	↑ ↑	I 0		No
	Hou et al. (2015) [#]	Taiwan	2012–2013	7–15	Cross-sectional	270	Urine	Nonylphenol	↑	↑	↑	9	I	No
	Lam et al. (2014)	Russia	2003–2005	8–9	Longitudinal cohort	350	Serum	p,p'-DDE HCB β-HCH	↑ ↓ ↑	↑ ↑ ↑	↑ ↑ ↑	9 0		No
	Ye et al. (2017) [#]	China	2014–2015	9–16	Cross-sectional	463	Urine	3-PBA	↓	↑	↑	8	I	No
Triclosan	Harley et al. (2019)	USA	1999–2000	9	Longitudinal cohort	139	Urine	TCS	↓	↑	↑	I 0		No

↑ Later onset; ↓ Earlier onset; ↔ none detected

[#]Self-reported puberty outcome rather than clinical examination

[§]CR: Completeness of reporting ≥ 8 was considered sufficient; Bias: I = potential risk of bias, 0 = lower risk of bias

^{§§}Measured postnatally in maternal biospecimen, but indicated as a proxy for prenatal exposure

*p-value < 0.05

(GS, TV, P): Included in meta-analysis for genital stage (GS); testicular volume (TV); pubarche (P)

**Sums of selected phthalate metabolites and substitutes:

ΣDEHPm: Molar sum of MEHP, 5OH-MEHP, 5oxo-MEHP, 5oxo-MEHP and 2oxo-MEHP expressed as DEHP

ΣDINPm: Molar sum of MINP, OH-MINP, 5oxo-MINP and 2oxo-MINP expressed as DINP

Low/high MWP: Low/high molecular weight phthalates

EDC, endocrine disrupting chemical.

et al., 2015; Burns et al., 2016; Namulanda et al., 2016, 2017; Harley et al., 2017, 2019; Kasper-Sonnenberg et al., 2017; Miao et al., 2017; Wang et al., 2017; Ye et al., 2017a,b; Berger et al., 2018; Binder et al., 2018; Attfield et al., 2019; Ernst et al., 2019; Cathey et al., 2020; Dobraca et al., 2020; Howland et al., 2020; Huang et al., 2020) that were based on 26 study populations reporting associations between prenatal and/or postnatal exposures to xenobiotic EDCs and timing of puberty outcomes for girls and boys: menarche (n = 31, self-report recall/self-report prospective diary: 30/1), thelarche (n = 25, clinical visual assessment/clinical assessment with breast palpitation/self-report: 1/18/6), pubarche in girls (n = 23, clinical assessment/self-report: 19/4), genital stage (n = 17, clinical assessment/self-report: 14/3), testicular volume (n = 15, orchidometer/self-report: 14/1) and pubarche in boys (n = 20, clinical assessment/self-report: 16/4). All 52 publications were used in the qualitative trend synthesis and 23 publications were eligible for meta-analysis.

Characteristics of the 52 publications in the systematic review are presented in Tables II and III. Overall, almost twice as many publications investigated puberty in girls (n = 41) compared with boys (n = 23), and most publications investigated more than one pubertal milestone as the outcome (n = 33). Assessment of associations according to postnatal EDC exposures (n = 36) was more prevalent compared with prenatal exposures to EDCs (n = 20) and only four publications assessed separate effects in both prenatal and postnatal exposure periods. Publications on prenatal exposure mainly assessed exposure during the second and/or third trimester (n = 9), while other publications collected samples prior to pregnancy (n = 4), in first trimester only (n = 1) or did not specify the time of collection within pregnancy (n = 3). Three publications collected maternal biospecimens after pregnancy: one at the time of delivery, one during breast feeding and one at the time of physical examinations of the child used as proxy for prenatal exposure. Publications on postnatal exposure mainly collected samples of biospecimen in the child prior to age 15 years (n = 25), but the overall age at exposure ranged from 5 to 19 years. In the majority of publications (n = 42), single spot biosamples were used to assess exposures in the study population, while in the remaining publications, biosamples were collected multiple times (up to six times) in a given period (data not shown). Of the 11 EDC compound groups investigated, most publications addressed exposure to phthalates (n = 19), polychlorinated phenols/pesticides (n = 18) and BPA (n = 12). The investigations predominantly used a longitudinal cohort design (60%) followed by a cross-sectional study design (35%), and most investigations were conducted on European (33%) and American (35%) study populations. Many individual publications were based on the same study populations, including The Breast Cancer and The Environment Research Program (BCERP) cohort (n = 7), Early Life Exposure in Mexico to Environmental Toxicants (ELEMENT) (n = 5), National Health and Nutrition Examination Survey (NHANES) (n = 4) and The Russian Children's Study (n = 4) (Supplementary Table SIV). Almost all included publications were published within the last 15 years (90%).

The completeness of reporting was generally high and only five publications were evaluated to have a low completeness of reporting. However, according to the predefined criteria, 50% of the included publications had a potential risk of bias primarily due to the potential for selection and information bias (Fig. 2).

Qualitative trend synthesis (n = 52 publications)

The qualitative trend synthesis provided data on 103 combinations of associations between prenatal or postnatal exposure to EDC compound groups and puberty outcomes. Selected trends (Tables II and III) by publication and Supplementary Table SIII by individual EDC in Supplementary data) are described below with a focus on the most investigated exposures and significant trends.

Menarche, thelarche and pubarche in girls

Three publications reported on prenatal exposure to BPA and ages at menarche, thelarche and pubarche, of which two were based on the same study population (Watkins et al., 2014, 2017b). These two publications indicated that exposure to BPA was associated with earlier thelarche and pubarche but later menarche, while the third publication with the largest study population (n = 537; Berger et al., 2018) consistently showed later onset for all three puberty outcomes (statistically significant for thelarche). Prenatal exposure to phthalates showed an overall trend toward earlier menarche, thelarche and pubarche, albeit the larger study by Berger et al. (2018) was an outlier and reported statistically significant associations with later onsets of puberty. Considering individual phthalate metabolites, prenatal exposure to MEP was consistently associated with earlier onset for all three outcomes. Prenatal exposure to polychlorinated phenols/pesticides showed a predominant trend toward earlier onset across all three puberty outcomes with several statistically significant associations for menarche.

Publications on postnatal exposure to BPA showed a relatively consistent trend toward later menarche and thelarche, but no clear trend for pubarche. Of the 12 publications on postnatal exposure to phthalates and pubertal timing in girls providing over 250 individual associations, no overall trend was observed with inconsistent directions of findings. Considering individual phthalate metabolites, postnatal exposure to MBzP, MEHP and \sum DEHPm seemed to support associations with earlier thelarche and later pubarche. Similarly, three publications on postnatal exposure to triclosan (Wolff et al., 2010, 2015; Harley et al., 2019) showed a clear trend toward earlier thelarche but tended toward later pubarche. Postnatal exposure to parabens showed a consistent trend toward earlier thelarche and pubarche, but less consistent findings for menarche. Postnatal exposure to polychlorinated phenols/pesticides did not show any clear trends for either menarche, thelarche or pubarche.

Genital stage, testicular volume and pubarche in boys

For puberty outcomes in boys, prenatal exposure to BPA showed a small trend toward earlier genital development, testicular enlargement and pubarche. Prenatal exposure to phthalates showed no trend for genital development and pubarche, but a predominant trend toward earlier onset was observed for testicular enlargement, although the most recent study by Cathey et al. (2020) indicated later onset.

Postnatal exposure to phthalates showed a weak trend toward later pubarche and earlier testicular enlargement, but only one publication investigated genital development, indicating earlier genital growth. Based on three publications, postnatal exposure to PCBs was associated with earlier onset across all puberty outcomes for boys. Publications on postnatal exposure to polychlorinated phenols/

pesticides showed a trend toward earlier onset of genital growth and pubarche, but no consistent trend for testicular enlargement.

Across all puberty outcomes in both girls and boys, prenatal exposures to UV filters and triclosan were based on a single study each and postnatal exposure to UV filters showed no clear overall trends. Exposures to dioxins and PAHs were generally investigated by too few studies to draw any overall conclusions.

Meta-analysis (n = 23 publications)

The meta-analysis of 23 publications enabled 18 summary risk estimates (prenatal/postnatal: 6/12) on pubertal onset in the main analysis (Strategy 1) following exposure to BPA, flame retardants, phthalates, polychlorinated phenols/pesticides, triclosan and UV filters. We observed varying heterogeneity in the estimates across all puberty outcomes and exposures (I^2 of 0–88%). Selected forest plots of the main analysis are included and presented in the manuscript in Figs 3 and 4, whereas the remaining forest plots are provided in Supplementary Figs S1 and S2.

Specific compound groups of EDCs (main analysis, Strategy 1)

Menarche. Twelve publications provided seven summary risk estimates for age at menarche following exposures to BPA, flame retardants, polychlorinated phenols/pesticides, phthalates, triclosan and UV filters. For prenatal exposures, only the summary estimate for prenatal exposure to phthalates was obtained and was close to unity (RR 1.01, 95% CI: 1.00–1.02; Fig. 3). Postnatal exposure to BPA showed a statistically non-significant association with later age at menarche (RR 0.95, 95% CI: 0.78–1.16) (Fig. 4A). But postnatal exposure to polychlorinated phenols/pesticides (Fig. 4B), flame retardants (Fig. 4C) and UV filters (Fig. 4D) was marginally associated with earlier age at menarche (RR 1.09, 95% CI: 0.99–1.19; RR 1.30, 95% CI: 0.94–1.79 and RR

1.04, 95% CI: 0.96–1.13, respectively). No associations were observed for postnatal exposure to phthalates (Fig. 4E) or triclosan (Fig. 4F), both with summary estimates of 1.00.

Thelarche. Seven publications provided three summary estimates for thelarche following exposures to BPA and phthalates. Prenatal exposure to phthalates showed no difference (RR 1.00, 95% CI: 0.99–1.01; Supplementary Fig. S1a). Postnatal exposure to BPA was associated with a statistically non-significant risk of later thelarche (RR 0.97, 95% CI: 0.87–1.09; Fig. 4G). No association was observed for postnatal exposure to phthalates and thelarche (RR 1.01, 95% CI: 1.00–1.02; Fig. 4H).

Pubarche in girls. Six publications provided three summary estimates for pubarche in girls following exposures to BPA, flame retardants and phthalates. No association was observed for prenatal exposure to phthalates and pubarche in girls (RR 1.01, 95% CI: 1.00–1.02) (Supplementary Fig. S1b). Postnatal exposures to BPA and phthalates were not associated with onset of pubarche (RR 1.01, 95% CI: 0.88–1.15 and RR 1.01, 95% CI: 1.00–1.02, respectively; Supplementary Fig. S2a and b).

Genital stage, testicular volume and pubarche in boys. Three publications provided five summary estimates for genital stage ($n = 1$), testicular volume ($n = 2$) and pubarche ($n = 2$) only following exposure to phthalates. The estimated meta-associations for boys were few and only based on a maximum of two different publications. Prenatal exposure to phthalates was associated with later genital development and pubarche onset (RR 0.98, 95% CI: 0.96–1.00 and RR 0.91, 95% CI: 0.84–0.98, respectively), but no association was observed for testicular enlargement (Supplementary Fig. S1c–e). Postnatal exposure to phthalates showed no association with testicular enlargement or pubarche in boys (Supplementary Fig. S2c and d).




Figure 3. Summary estimates of the meta-analysis: associations between prenatal exposure to phthalates and their metabolites and menarche.

Persistent and non-persistent EDC compounds (Strategy 2)

Puberty outcomes in girls. Grouping EDCs into persistent and non-persistent compounds resulted in summary estimates for prenatal exposures that were based on relatively few publications. Prenatal

exposures to non-persistent compounds were based on the same two publications by [Watkins et al. \(2014, 2017b\)](#) and showed essentially no association across all puberty outcomes in girls ([Supplementary Fig. S1f–h](#)). Prenatal exposure to persistent compounds was associated




Figure 4. Summary estimates of the meta-analysis (Strategy 1): postnatal exposures and puberty outcomes in girls. (A) Associations between postnatal exposure to bisphenol A and menarche. **(B)** Associations between postnatal exposure to polychlorinated phenols and pesticides and menarche. **(C)** Associations between postnatal exposure to flame retardants and menarche. **(D)** Associations between postnatal exposure to benzophenones and other UV filters and menarche. **(E)** Associations between postnatal exposure to phthalates and their metabolites and menarche. **(F)** Associations between postnatal exposure to triclosan and menarche. **(G)** Associations between postnatal exposure to bisphenol A and thelarche. **(H)** associations between postnatal exposure to phthalates and their metabolites and thelarche.




Figure 4. (Continued)

with later age at menarche (0.73, 95% CI: 0.63–0.84; [Supplementary Fig. S1i](#)).

Postnatal exposure to non-persistent compounds was associated with slightly earlier thelarche (RR 1.02, 95% CI: 0.99–1.05), but no difference for menarche and pubarche (RR 1.00, 95% CI: 1.00–1.01 and RR 1.01, 95% CI: 1.00–1.02) ([Supplementary Fig. S2e–g](#)). Postnatal exposure to persistent compounds showed statistically non-significant associations with earlier age at menarche (RR 1.09, 95% CI: 0.88–1.37) and later age at pubarche (RR 0.96, 95% CI: 0.80–1.14), but a significant association with later thelarche (odds ratio 0.73, 95% CI: 0.59–0.90) ([Supplementary Fig. S2h–j](#)).

Puberty outcomes in boys. Meta-associations for genital development, testicular enlargement and pubarche in boys following prenatal or postnatal exposures to non-persistent compounds were largely based on the same individual estimates as for exposure to phthalates and reflected the same subsequent summary estimates ([Supplementary Fig. S1j–l](#) and [S2k–l](#)). No summary estimates were enabled for exposures to persistent compounds.

Banned and suspicious EDC compounds (Strategy 3)

In the analysis of banned and suspicious exposure compounds, prenatal or postnatal exposures to EDCs were not associated with pubertal onset, with most of the non-statistically significant summary estimates ranging between RR 0.99 and 1.01 (data not shown).

Discussion

This systematic review pooled available evidence on the association between prenatal or postnatal exposures to xenobiotic EDCs and timing of pubertal onset in girls and boys. Although the findings of this review provide some evidence on the effect of exposure to EDCs on

pubertal onset, the available data on specific exposure-outcome associations are limited and thus preclude strong concluding statements. The qualitative trend synthesis suggested that postnatal exposure to phthalates may be associated with an earlier thelarche and later pubarche. However, in the subsequent stringent meta-analysis, no clear overall association of EDCs on puberty development in girls and boys was evident across all outcomes. Based on current evidence, we were not able to identify pre- or postnatal windows of exposure as particularly critical and susceptible for effects of EDCs.

Overall findings of the review

Qualitative trend synthesis

Many of the identified trends in the qualitative trend synthesis of 52 publications were often subject to inconsistent findings within specific exposure-outcome associations and/or sometimes contradicting directions of findings across puberty outcomes for the same compound group. For a few specific exposure-outcome associations, the trend synthesis was able to provide an overall indication of an association, such as largely consistent trends toward later menarche and later thelarche following postnatal exposure to BPA. In the trend synthesis, we were also able to identify an association between postnatal exposure to phthalates (the compound group most investigated) and earlier thelarche but later pubarche, which could not be confirmed in the meta-analysis, potentially due to a lower number of included publications in our meta-analysis. These findings may also in part reflect different hormonal actions of thelarche (an estrogen-driven process) and pubarche (an androgen-driven process). However, the individual findings for prenatal exposure to phthalates indicated an overall direction toward earlier onset of all puberty outcomes in girls (menarche, thelarche, pubarche), although this trend was based on few publications and was relatively weak. For puberty in boys, the trend synthesis was not able

to identify any clear trends, probably as these were based on very few individual exposure-outcome associations. Despite some suggestive associations in the qualitative trend synthesis, many identified trends were not robust, and the overall effect of specific exposure compounds remained largely inconclusive.

Meta-analysis

The summary estimates from the meta-analysis of 23 publications provided scarce evidence on altered age of pubertal onset following prenatal or postnatal exposures to EDCs, largely due to difficulties in obtaining actual meta-associations based on sufficient data. Most exposure-outcome associations showed no overall effect, and summary risk estimates with indication of an association showed relatively small changes in pubertal onset with limited predictive power. On balance, the meta-associations provided less evidence for the included compounds with more individual associations, i.e. summary estimates on exposure to phthalates were based on most individual estimates but the resulting summary estimates generally indicated no associations. Applying different strategies did not provide additional perspectives on the overall findings. A few statistically significant associations for persistent compounds suggested later onset of puberty in girls, but given the many analyses performed, these individual associations may be a result of chance findings.

Quality, limitations and comparability of the included publications

The specific reporting quality of each of the 52 included publications was considered sufficiently high, though with a noteworthy potential risk of selection and information bias. In a broader sense, the overall research quality within the publication landscape was subject to several methodological challenges and inconsistencies. Most publications ascertained the respective puberty outcomes in a homogenous way using standardized clinical assessments, including visualization and palpation by trained clinicians, by standards of the Tanner staging of development. However, definitions of pubertal onset were surprisingly different and not all publications defined onset of puberty as having reached Tanner Stage 2. Regarding the use of self-reported data for age of menarche, some epidemiological publications prospectively followed and assessed menarche in children using a diary, whereas others solely relied on a one-time retrospective assessment with risk of recall bias and potential outcome misclassification. Standardized clinical definitions of pubertal onset across all puberty outcomes are therefore needed for future research to allow for formal comparison of evidence. Exposure assessments were relatively comparable across publications as we only included publications with reliable exposure biomarkers, but the specific levels of exposure were still assessed by different sample handling practices such as storage containers and storage time (historical cohorts), which may lead to EDC degradation or sample contamination. Importantly, timing of exposure assessment also varied significantly between publications and the specific exposure period may impact the association, a possible concern for *in utero* exposure during susceptible periods of fetal development or different windows throughout childhood. The predominant use of single spot biosamples during pregnancy and in childhood challenges interpretation of associations with full certainty and a risk of exposure misclassification is inevitable for non-persistent compounds, and potentially also for persistent compounds, especially depending on the specific chemical and source of exposure (diet, household products, cosmetics, etc.; Bloom *et al.*,

2007). Across all publications, there were generally important differences in methodological choices, especially regarding study design, sample size, statistical analyses, adjustment for creatinine, specific gravity and serum lipids and methods of log-transforming EDC exposure levels. Although most publications used a longitudinal cohort study design, a significant number of publications were cross-sectional despite the predictive limitations in assessing exposures at the same time as outcome with the risk of reverse causality. All these observed inconsistencies in methodological approaches may subsequently preclude any strong conclusions on the impact of EDCs on onset of puberty.

Critical windows of fetal development

Pregnancy is a critical period of development, during which the fetus is susceptible to harmful effects of exposures. Healthy fetal development largely depends on hormonal regulation and any disturbances of hormones could ultimately affect growth and development parameters. Therefore, we expected that prenatal exposures to EDCs might interfere with development of fetal endocrine systems and represent a critical window of exposure with regard to timing of pubertal onset. However, based on this review, we were unable to recognize prenatal exposures as particularly critical for effects of EDCs and evidence was too scarce to distinguish different windows of exposure during pregnancy.

Combined mixtures of EDC exposures

International biomonitoring data from the past four decades provide unequivocal documentation of human exposure to a mixture of xenobiotic substances, including persistent and non-persistent EDCs. There is increasing interest in ascertaining joint effects of chemicals, as chemical exposures rarely occur in isolation and the potential effects on pubertal perturbations likely depend on a mixture of exposures present in low concentrations rather than attributable to single EDC compounds. However, we were not able to evaluate the simultaneous exposures to different EDCs (the so-called cocktail effect; Kortenkamp, 2007, 2014), as all included publications considered one EDC compound at a time and we were unable to draw conclusions on the potentially additive or synergistic interactions between the investigated EDC compounds on the pubertal outcomes. In our meta-analyses, we attempted to address the combined joint effects within and between different classes of EDC compounds by pooling estimates according to various strategies (according to compound groups of EDCs, persistency and chemicals on the SVHC lists). We acknowledge that these analyses may be over-simplified, as the different EDCs vary in their biological modes of actions and potencies, but biological actions are not fully elucidated in the existing literature and ascertaining this with full certainty was beyond the scope of this systematic review.

Biological plausibility

Numerous studies have explored exposures to EDCs in relation to pubertal onset, but mostly focused on individual compounds and their association with timing of specific pubertal outcomes. In line with our findings from the trend synthesis, we previously reported that childhood excretion (biomarker of exposure due to correlation) of high levels of urinary phthalate metabolites was associated with a marked delay in the development of pubarche among girls (Frederiksen *et al.*, 2012). Similar findings were confirmed shortly hereafter in a US study (Wolff *et al.*, 2014), whereas others did not demonstrate an

association. A delay in pubarche may potentially be explained by the anti-androgenic properties of phthalates and result in later development of pubarche (Axelstad et al., 2018). Despite a plausible biological mechanism, the varying anti-androgenic properties of individual phthalate metabolites were not accounted for in the respective study and the potential underlying mechanism remains largely speculative. Importantly, neither of these two mentioned studies reported their data in a way that allowed for their inclusion in our meta-analysis.

Strengths and limitations of the review

We have rigorously evaluated the evidence on prenatal and postnatal exposures using biomarkers to EDC compounds and pubertal onset in girls and boys. To our knowledge, this review includes all published studies in English (prior to 1 February 2021) fulfilling predefined criteria using a systematic and transparent search of the literature. Unlike previous systematic reviews, we included both a qualitative trend synthesis and a meta-analysis with summary risk estimates applying different approaches for exposure groupings.

However, an important limitation of our systematic review is the rather limited number of available studies for investigating specific exposure (by compound)-outcome associations and some of the summary risk estimates in the meta-analysis lack sufficient data to allow for reliable meta-associations. Given the limited data, we were compelled to investigate exposure in childhood under two broadly defined exposure categories, within fetal life and during childhood (after birth). We were not able to investigate finer windows of exposure for which there may be differences in susceptibility and effects of exposure, e.g. early childhood versus pre-puberty. Another inherent limitation is the methodological heterogeneity among studies in study design, statistical analyses, definitions of puberty etc. that resulted in a substantial but necessary exclusion of publications in the meta-analysis to enable comparison of individual estimates. Even though these limitations in comparing studies were unavoidable, interpretation of the meta-associations on the impact of EDCs should be made with caution.

As previously discussed, our review is also limited by only assessing individual effects of EDC compounds and therefore not including possible synergies and mixture effects, even though a combined exposure to multiple EDCs may in fact represent the underlying explanation to any observed association between EDCs and timing of pubertal onset.

Conclusions and wider implications

In this large systematic review with a stringent meta-analysis, we did not find consistent evidence for associations between prenatal or postnatal exposures to xenobiotic EDCs and changes in pubertal timing in girls and boys. According to statistically significant associations in the qualitative trend synthesis, postnatal exposure to phthalates may be associated with earlier thelarche and later pubarche, consistent with their anti-androgenic properties. Only a few specific exposure-outcome associations were identified in the meta-analysis but with limited predictive power. We were not able to identify specific pre- or postnatal windows of exposure as particularly critical and susceptible for effects of EDCs. Current evidence is subject to several methodological challenges and inconsistencies and the evidence on specific exposure-outcome associations remains too scarce to firmly confirm EDC exposure as a risk factor for change in age of pubertal onset. Adding to the complexity of this research field, virtually all children are

exposed to a multitude of varying EDCs throughout their fetal and postnatal lifetime and these exposures often correlate, which makes identifying individual EDC contributions to altered timing of pubertal onset difficult to ascertain. To create a more uniform foundation for future comparison of evidence and to strengthen pooled studies, we recommend the use of more standardized approaches in the choice of statistical analyses, with transformation of non-normal distributed EDC exposures, and in the definitions and assessments of puberty outcomes. The impact of mixtures of EDC exposures on the association also remains unestablished and would be valuable to elucidate for prenatal and postnatal windows of exposure. Future large, longitudinal epidemiological studies are needed to clarify the impact of EDCs on pubertal timing, as some EDC exposures are increasing and have potential health implications for young children.

Supplementary data

Supplementary data are available at *Human Reproduction Update* online.

Data availability

All data are incorporated into the article and its online [supplementary data](#).

Acknowledgements

We are extremely grateful to the librarian, Susie Rimborg, at the University of Copenhagen library for her assistance with the systematic literature search protocol in the PubMed database.

Authors' roles

A.J. and E.V.B. designed the concept of the study and supervised the study. T.K. performed the systematic literature search and screening of publications with support from E.V.B. T.K. and C.S.O. evaluated the quality of the included studies. T.K. and C.S.U. drafted the tables with support from E.V.B., while L.S.G. validated the content of the final table versions. C.S.U. and T.K. performed the qualitative analyses with support from L.S.G. and E.V.B. Y.H.L. performed the statistical analyses. A.M.A. and H.F. provided EDC expertise and strategies for grouping exposure compounds. B.A.C. provided guidance on statistical methods and R.H. provided expertise in relation to pubertal onset. C.S.U. drafted the manuscript with support from T.K. and E.V.B. All authors contributed to manuscript revisions and the final draft of the manuscript.

Funding

The salaries of A.J., E.V.B., C.S.U. and A.M.A. were partially supported by a grant from the National Institute of Health (grant no. IR01CA236816-01A1). E.V.B. was also partially supported by grants from the Danish Health Foundation (Helsefonden, grant no. 18-B-

0016, Aase & Ejnar Danielsen, grant no. 10-002122, Svend Andersen Fonden, grant no 81A-01 and Familien Erichsens Mindefond, grant no. 6000073). T.K. was also supported by the Danish Health Foundation (Helsefonden, grant no. 19-B-0077). C.S.U. was also partially supported by a grant from the Danish Cancer Research Association (Kræftens Bekæmpelse, grant no. R204-A12636). L.S.G. was supported by a grant from the Danish Environmental Protection Agency (Miljøstyrelsen, grant no. MST-611-00012). BC was funded by a grant from the National Institutes of Health (NIH) (grant no. ES000002).

Conflict of interest

None declared.

References

- Akslglaede L, Sørensen K, Petersen JH, Skakkebaek NE, Juul A. Recent decline in age at breast development: the Copenhagen puberty study. *Pediatrics* 2009;**123**:e932–e939.
- Attfield KR, Pinney SM, Sjödin A, Voss RW, Greenspan LC, Biro FM, Hiatt RA, Kushi LH, Windham GC. Longitudinal study of age of menarche in association with childhood concentrations of persistent organic pollutants. *Environ Res* 2019;**176**:108551.
- Axelstad M, Hass U, Scholze M, Christiansen S, Kortenkamp A, Boberg J. EDC IMPACT: Reduced sperm counts in rats exposed to human relevant mixtures of endocrine disrupters. *Endocr Connect* 2018;**7**:139–148.
- Berger K, Eskenazi B, Kogut K, Parra K, Lustig RH, Greenspan LC, Holland N, Calafat AM, Ye X, Harley KG. Association of prenatal urinary concentrations of phthalates and bisphenol A and pubertal timing in boys and girls. *Environ Health Perspect* 2018;**126**:097004–097009.
- Binder AM, Corvalan C, Calafat AM, Ye X, Mericq V, Pereira A, Michels KB. Childhood and adolescent phenol and phthalate exposure and the age of menarche in Latina girls. *Environ Heal A Glob Access Sci Source* 2018;**17**:1–11.
- Blanck HM, Marcus M, Tolbert PE, Rubin C, Henderson AK, Hertzberg VS, Zhang RH, Cameron L. Age at menarche and tanner stage in girls exposed in utero and postnatally to polybrominated biphenyl. *Epidemiology* 2000;**11**:641–647.
- Bloom MS, Buck Louis GM, Schisterman EF, Liu A, Kostyniak PJ. Maternal serum polychlorinated biphenyl concentrations across critical windows of human development. *Environ Health Perspect* 2007;**115**:1320–1324.
- Bonilla C, Lewis SJ, Martin RM, Donovan JL, Hamdy FC, Neal DE, Eeles R, Easton D, Kote-Jarai Z, Al Olama AA et al.; PRACTICAL Consortium. Pubertal development and prostate cancer risk: Mendelian randomization study in a population-based cohort. *BMC Med* 2016;**14**:66.
- Bonzini M, Coggon D, Palmer KT. Risk of prematurity, low birth-weight and pre-eclampsia in relation to working hours and physical activities: a systematic review. *Occup Environ Med* 2007;**64**:228–243.
- Bundesumwelt U, Kem M. European workshop on the impact of endocrine disrupters on human health and wildlife. Report of Proceedings from a Workshop Held in Weybridge, UK, 1996. In European Commission, DGXII, Brussels, Belgium.
- Burns JS, Lee MM, Williams PL, Korrick SA, Sergeyev O, Lam T, Revich B, Hauser R. Associations of peripubertal serum dioxin and polychlorinated biphenyl concentrations with pubertal timing among Russian boys. *Environ Health Perspect* 2016;**124**:1801–1807.
- Buttke DE, Sircar K, Martin C. Exposures to endocrine-disrupting chemicals and age of menarche in adolescent girls in NHANES (2003–2008). *Environ Health Perspect* 2012;**120**:1613–1618.
- Castiello F, Freire C. Exposure to non-persistent pesticides and puberty timing: a systematic review of the epidemiological evidence. *Eur J Endocrinol* 2021;**184**:733–749.
- Cathey A, Watkins DJ, Sánchez BN, Tamayo-Ortiz M, Solano-Gonzalez M, Torres-Olascoaga L, Téllez-Rojo MM, Peterson KE, Meeker JD. Onset and tempo of sexual maturation is differentially associated with gestational phthalate exposure between boys and girls in a Mexico City birth cohort. *Environ Int* 2020;**136**:105469.
- Chen A, Chung E, DeFranco EA, Pinney SM, Dietrich KN. Serum PBDEs and age at menarche in adolescent girls: analysis of the National Health and Nutrition Examination Survey 2003–2004. *Environ Res* 2011;**111**:831–837.
- Christensen KY, Maisonet M, Rubin C, Holmes A, Calafat AM, Kato K, Flanders WD, Heron J, McGeehin MA, Marcus M. Exposure to polyfluoroalkyl chemicals during pregnancy is not associated with offspring age at menarche in a contemporary British cohort. *Environ Int* 2011;**37**:129–135.
- Day FR, Elks CE, Murray A, Ong KK, Perry JRB. Puberty timing associated with diabetes, cardiovascular disease and also diverse health outcomes in men and women: the UK Biobank study. *Sci Rep* 2015;**5**:11208.
- Day FR, Thompson DJ, Helgason H, Chasman DI, Finucane H, Sulem P, Ruth KS, Whalen S, Sarkar AK, Albrecht E et al.; The LifeLines Cohort Study. Genomic analyses identify hundreds of variants associated with age at menarche and support a role for puberty timing in cancer risk. *Nat Genet* 2017;**49**:834–841.
- Denham M, Schell LM, Deane G, Gallo MV, Ravenscroft J, DeCaprio AP; The Akwesasne Task Force on the Environment. Relationship of lead, mercury, mirex, dichlorodiphenyldichloroethylene, hexachlorobenzene, and polychlorinated biphenyls to timing of menarche among Akwesasne Mohawk girls. *Pediatrics* 2005;**115**:e127–e134.
- Dobrca D, Laurent CA, Greenspan LC, Hiatt RA, Sjödin A, Kushi LH, Windham GC. Urinary polycyclic aromatic hydrocarbons in relation to anthropometric measures and pubertal development in a cohort of Northern California girls. *Environ Epidemiol* 2020;**4**:e0102.
- Domínguez-Romero E, Scheringer M. A review of phthalate pharmacokinetics in human and rat: what factors drive phthalate distribution and partitioning? *Drug Metab Rev* 2019;**51**:314–329.
- Elks CE, Ong KK, Scott RA, Van Der Schouw YT, Brand JS, Wark PA, Amiano P, Balkau B, Barricarte A, Boeing H et al.; The InterAct Consortium. Age at menarche and type 2 diabetes risk: The EPIC-InterAct study. *Diabetes Care* 2013;**36**:3526–3534.
- Ernst A, Brix N, Lauridsen LLB, Olsen J, Parnert ET, Liew Z, Olsen LH, Ramlau-Hansen CH. Exposure to perfluoroalkyl substances during fetal life and pubertal development in boys and girls from

- the Danish national birth cohort. *Environ Health Perspect* 2019; **127**:017004.
- European Chemicals Agency. Registrerede stoffer - ECHA. 2021a. <https://echa.europa.eu/da/information-on-chemicals/registered-substances> (11 August 2021, date last accessed).
- European Chemicals Agency. *The Candidate List of Substances of Very High Concern now Contains 219 Chemicals that may Harm People or the Environment*. 2021b. <https://echa.europa.eu/da/-/candidate-list-updated-with-eight-hazardous-chemicals> (18 August 2021, date last accessed).
- Ferguson KK, Peterson KE, Lee JM, Mercado-García A, Blank-Goldenberg C, Téllez-Rojo MM, Meeker JD. Prenatal and peripubertal phthalates and bisphenol A in relation to sex hormones and puberty in boys. *Reprod Toxicol* 2014; **47**:70–76.
- Frederiksen H, Sørensen K, Mouritsen A, Aksglaede L, Hagen CP, Petersen JH, Skakkebaek NE, Andersson AM, Juul A. High urinary phthalate concentration associated with delayed pubarche in girls. *Int J Androl* 2012; **35**:216–226.
- Grandjean P, Grønlund C, Kjær IM, Jensen TK, Sørensen N, Andersson AM, Juul A, Skakkebaek NE, Budtz-Jørgensen E, Weihe P. Reproductive hormone profile and pubertal development in 14-year-old boys prenatally exposed to polychlorinated biphenyls. *Reprod Toxicol* 2012; **34**:498–503.
- Hagan H, Pouget ER, Des Jarlais DC. A systematic review and meta-analysis of interventions to prevent Hepatitis C virus infection in people who inject drugs. *J Infect Dis* 2011; **204**:74–83.
- Hamlat EJ, Stange JP, Abramson LY, Alloy LB. Early pubertal timing as a vulnerability to depression symptoms: differential effects of race and sex. *J Abnorm Child Psychol* 2014; **42**:527–538.
- Harley KG, Berger KP, Kogut K, Parra K, Lustig RH, Greenspan LC, Calafat AM, Ye X, Eskenazi B. Association of phthalates, parabens and phenols found in personal care products with pubertal timing in girls and boys. *Hum Reprod* 2019; **34**:109–117.
- Harley KG, Rauch SA, Chevrier J, Kogut K, Parra KL, Trujillo C, Lustig RH, Greenspan LC, Sjödin A, Bradman A et al. Association of prenatal and childhood PBDE exposure with timing of puberty in boys and girls. *Environ Int* 2017; **100**:132–138.
- Hartemink N, Boshuizen HC, Nagelkerke NJD, Jacobs MAM, van Houwelingen HC. Combining risk estimates from observational studies with different exposure cutpoints: a meta-analysis on body mass index and diabetes type 2. *Am J Epidemiol* 2006; **163**:1042–1052.
- Herman-Giddens ME, Slora EJ, Wasserman RC, Bourdony CJ, Bhapkar MV, Koch GG, Hasemeier CM. Secondary sexual characteristics and menses in young girls seen in office practice: A study from the pediatric research in office settings network. *Pediatrics* 1997; **99**:505–512.
- Herman-Giddens ME, Wang L, Koch G. Secondary sexual characteristics in boys: Estimates from the National Health and Nutrition Examination Survey III, 1988–1994. *Arch Pediatr Adolesc Med* 2001; **155**:1022–1028.
- Den Hond E, Dhooze W, Bruckers L, Schoeters G, Nelen V, Van De Mieroop E, Koppen G, Bilau M, Schroyen C, Keune H et al. Internal exposure to pollutants and sexual maturation in Flemish adolescents. *J Expo Sci Environ Epidemiol* 2011; **21**:224–233.
- Hou JW, Lin CL, Tsai YA, Chang CH, Liao KW, Yu CJ, Yang W, Lee MJ, Huang PC, Sun CW et al. The effects of phthalate and nonylphenol exposure on body size and secondary sexual characteristics during puberty. *Int J Hyg Environ Health* 2015; **218**:603–615.
- Howland RE, Deziel NC, Bentley GR, Booth M, Choudhury OA, Hofmann JN, Hoover RN, Katki HA, Trabert B, Fox SD et al. Assessing endogenous and exogenous hormone exposures and breast development in a migrant study of Bangladeshi and British girls. *Int J Environ Res Public Health* 2020; **17**:1185.
- Huang Y, Wang P, Law JCF, Zhao Y, Wei Q, Zhou Y, Zhang Y, Shi H, Leung KSY. Organic UV filter exposure and pubertal development: a prospective follow-up study of urban Chinese adolescents. *Environ Int* 2020; **143**:105961.
- Humblet O, Williams PL, Korrick SA, Sergeyev O, Emond C, Birnbaum LS, Burns JS, Altshul L, Patterson DG, Turner WE et al. Dioxin and polychlorinated biphenyl concentrations in mother's serum and the timing of pubertal onset in sons. *Epidemiology* 2011; **22**:827–835.
- Kaltiala-Heino R, Marttunen M, Aivi Rantanen P, Rimpel M. Early puberty is associated with mental health problems in middle adolescence. *Soc Sci Med* 2003; **57**:1055–1064.
- Kaplowitz PB, Oberfield SE. Reexamination of the age limit for defining when puberty is precocious in girls in the United States: implications for evaluation and treatment. *Pediatrics* 1999; **104**:936–941.
- Kasper-Sonnenberg M, Wittsiepe J, Wald K, Koch HM, Wilhelm M. Pre-pubertal exposure with phthalates and bisphenol A and pubertal development. *PLoS One* 2017; **12**:e0187922.
- Korrick SA, Lee MM, Williams PL, Sergeyev O, Burns JS, Patterson DG, Turner WE, Needham LL, Altshul L, Revich B et al. Dioxin exposure and age of pubertal onset among Russian boys. *Environ Health Perspect* 2011; **119**:1339–1344.
- Kortenkamp A. Ten years of mixing cocktails: a review of combination effects of endocrine-disrupting chemicals. *Environ Health Perspect* 2007; **115**:98–105.
- Kortenkamp A. Low dose mixture effects of endocrine disruptors and their implications for regulatory thresholds in chemical risk assessment. *Curr Opin Pharmacol* 2014; **19**:105–111.
- Lam T, Williams PL, Lee MM, Korrick SA, Birnbaum LS, Burns JS, Sergeyev O, Revich B, Altshul LM, Patterson DG et al. Prepubertal organochlorine pesticide concentrations and age of pubertal onset among Russian boys. *Environ Int* 2014; **73**:135–142.
- Lee JE, Jung HW, Lee YJ, Lee YA. Current evidence for the roles of early-life endocrine disruptors on pubertal timing in girls. *Ann Pediatr Endocrinol Metab* 2019; **24**:78–91.
- Leijds MM, Koppe JG, Olie K, Va Aalderen WMC, De Voogt P, Vulsma T, Westra M, ten Tusscher GW. Delayed initiation of breast development in girls with higher prenatal dioxin exposure; a longitudinal cohort study. *Chemosphere* 2008; **73**:999–1004.
- Lopez-Espinosa MJ, Fletcher T, Armstrong B, Genser B, Dhataria K, Mondal D, Ducatman A, Leonardi G. Association of perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) with age of puberty among children living near a chemical plant. *Environ Sci Technol* 2011; **45**:8160–8166.
- Lucaccioni L, Trevisani V, Marrozzini L, Bertoncelli N, Predieri B, Lugli L, Berardi A, Iughetti L. Endocrine-disrupting chemicals and their effects during female puberty: A review of current evidence. *Int J Mol Sci* 2020; **21**:2078.

- Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. *Arch Dis Child* 1969;**44**:291–303.
- McCarthy M. European Chemical Agency (ECHA) List of Pre-Registered Substances. 2011. <https://echa.europa.eu/da/information-on-chemicals/pre-registered-substances> (16 June 2021, date last accessed).
- McGuinn LA, Ghazarian AA, Joseph Su L, Ellison GL. Urinary bisphenol A and age at menarche among adolescent girls: evidence from NHANES 2003–2010. *Environ Res* 2015;**136**:381–386.
- Miao M, Wang Z, Liu X, Liang H, Zhou Z, Tan H, Yuan W, Li DK. Urinary bisphenol A and pubertal development in Chinese school-aged girls: a cross-sectional study. *Environ Heal A Glob Access Sci Source* 2017;**16**:1–7.
- Mouritsen A, Frederiksen H, Sorensen K, Aksglaede L, Hagen C, Skakkebaek NE, Main KM, Andersson AM, Juul A. Urinary phthalates from 168 girls and boys measured twice a year during a 5-year period: associations with adrenal androgen levels and puberty. *J Clin Endocrinol Metab* 2013;**98**:3755–3764.
- Namulanda G, Maisonet M, Taylor E, Flanders WD, Olson D, Sjodin A, Qualters JR, Vena J, Northstone K, Naeher L. In utero exposure to organochlorine pesticides and early menarche in the Avon longitudinal study of parents and children. *Environ Int* 2016;**94**:467–472.
- Namulanda G, Taylor E, Maisonet M, Boyd Barr D, Flanders WD, Olson D, Qualters JR, Vena J, Northstone K, Naeher L. In utero exposure to atrazine analytes and early menarche in the Avon longitudinal study of parents and children cohort. *Environ Res* 2017;**156**:420–425.
- Ouyang F, Perry MJ, Venners SA, Chen C, Wang B, Yang F, Fang Z, Zang T, Wang L, Xu X et al. Serum DDT, age at menarche, and abnormal menstrual cycle length. *Occup Environ Med* 2005;**62**:878–884.
- Parent AS, Teilmann G, Juul A, Skakkebaek NE, Toppari J, Bourguignon JP. The timing of normal puberty and the age limits of sexual precocity: variations around the world, secular trends, and changes after migration. *Endocr Rev* 2003;**24**:668–693.
- Prentice P, Viner RM. Pubertal timing and adult obesity and cardiometabolic risk in women and men: a systematic review and meta-analysis. *Int J Obes* 2013;**37**:1036–1043.
- Shamliyan T, Kane RL, Dickinson S. A systematic review of tools used to assess the quality of observational studies that examine incidence or prevalence and risk factors for diseases. *J Clin Epidemiol* 2010;**63**:1061–1070.
- Shi H, Cao Y, Shen Q, Zhao Y, Zhang Z, Zhang Y. Association between urinary phthalates and pubertal timing in Chinese adolescents. *J Epidemiol* 2015;**25**:574–582.
- Small CM, Terrell ML, Cameron LL, Wirth J, Monteilh CP, Marcus M. In utero exposure to a brominated flame retardant and male growth and development. *Int J Child Adolesc Health* 2009;**2**:11.
- Su PH, Chang CK, Lin CY, Chen HY, Liao PC, Hsiung CA, Chiang HC, Wang SL. Prenatal exposure to phthalate ester and pubertal development in a birth cohort in central Taiwan: A 12-year follow-up study. *Environ Res* 2015;**136**:324–330.
- Sun SS, Schubert CM, Chumlea WC, Roche AF, Kulin HE, Lee PA, Himes JH, Ryan AS. National estimates of the timing of sexual maturation and racial differences among US children. *Pediatrics* 2002;**110**:911–919.
- Sørensen K, Aksglaede L, Petersen JH, Juul A. Recent changes in pubertal timing in healthy Danish boys: associations with body mass index. *J Clin Endocrinol Metab* 2010;**95**:263–270.
- Sørensen K, Juul A, Christensen K, Skytthe A, Scheike T, Kold Jensen T. Birth size and age at menarche: a twin perspective. *Hum Reprod* 2013;**28**:2865–2871.
- Vasilu O, Muttineni J, Karmaus W. In utero exposure to organochlorines and age at menarche. *Hum Reprod* 2004;**19**:1506–1512.
- Wang Z, Li D, Miao M, Liang H, Chen J, Zhou Z, Wu C, Yuan W. Urine bisphenol A and pubertal development in boys. *Int J Hyg Environ Health* 2017;**220**:43–50.
- Warner M, Rauch S, Ames J, Mocarelli P, Brambilla P, Eskenazi B. Age at menarche in Seveso daughters exposed in utero to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Environ Epidemiol* 2020;**4**:e111.
- Warner M, Samuels S, Mocarelli P, Gerthoux PM, Needham L, Patterson DG, Eskenazi B. Serum dioxin concentrations and age at menarche. *Environ Health Perspect* 2004;**112**:1289–1292.
- Watkins DJ, Sánchez BN, Téllez-Rojo MM, Lee JM, Mercado-García A, Blank-Goldenberg C, Peterson KE, Meeker JD. Impact of phthalate and BPA exposure during in utero windows of susceptibility on reproductive hormones and sexual maturation in peripubertal males. *Environ Heal A Glob Access Sci Source* 2017a;**16**:1–10.
- Watkins DJ, Sánchez BN, Téllez-Rojo MM, Lee JM, Mercado-García A, Blank-Goldenberg C, Peterson KE, Meeker JD. Phthalate and bisphenol A exposure during in utero windows of susceptibility in relation to reproductive hormones and pubertal development in girls. *Environ Res* 2017b;**159**:143–151.
- Watkins DJ, Téllez-Rojo MM, Ferguson KK, Lee JM, Solano-Gonzalez M, Blank-Goldenberg C, Peterson KE, Meeker JD. In utero and peripubertal exposure to phthalates and BPA in relation to female sexual maturation. *Environ Res* 2014;**134**:233–241.
- Windham GC, Pinney SM, Voss RW, Sjodin A, Biro FM, Greenspan LC, Stewart S, Hiatt RA, Kushi LH. Brominated flame retardants and other persistent organohalogenated compounds in relation to timing of puberty in a longitudinal study of girls. *Environ Health Perspect* 2015;**123**:1046–1052.
- Wolff MS, Britton JA, Boguski L, Hochman S, Maloney N, Serra N, Liu Z, Berkowitz G, Larson S, Forman J. Environmental exposures and puberty in inner-city girls. *Environ Res* 2008;**107**:393–400.
- Wolff MS, Pajak A, Pinney SM, Windham GC, Galvez M, Rybak M, Silva MJ, Ye X, Calafat AM, Kushi LH et al. Associations of urinary phthalate and phenol biomarkers with menarche in a multiethnic cohort of young girls. *Reprod Toxicol* 2017;**67**:56–64.
- Wolff MS, Teitelbaum SL, McGovern K, Pinney SM, Windham GC, Galvez M, Pajak A, Rybak M, Calafat AM, Kushi LH et al.; Breast Cancer and Environment Research Program. Environmental phenols and pubertal development in girls. *Environ Int* 2015;**84**:174–180.
- Wolff MS, Teitelbaum SL, McGovern K, Windham GC, Pinney SM, Galvez M, Calafat AM, Kushi LH, Biro FM; on behalf of the Breast Cancer and Environment Research Program. Phthalate exposure and pubertal development in a longitudinal study of US girls. *Hum Reprod* 2014;**29**:1558–1566.
- Wolff MS, Teitelbaum SL, Pinney SM, Windham G, Liao L, Biro F, Kushi LH, Erdmann C, Hiatt RA, Rybak ME et al.; Breast Cancer and Environment Research Centers. Investigation of relationships between

- urinary biomarkers of phytoestrogens, phthalates, and phenols and pubertal stages in girls. *Environ Health Perspect* 2010;**118**:1039–1046.
- Wu T, Mendola P, Buck GM. Ethnic differences in the presence of secondary sex characteristics and menarche among US girls: The Third National Health and Nutrition Examination Survey, 1988–1994. *Pediatrics* 2002;**110**:752–757.
- Ye X, Pan W, Zhao S, Zhao Y, Zhu Y, Liu J, Liu W. Relationships of pyrethroid exposure with gonadotropin levels and pubertal development in Chinese boys. *Environ Sci Technol* 2017a;**51**:6379–6386.
- Ye X, Pan W, Zhao Y, Zhao S, Zhu Y, Liu W, Liu J. Association of pyrethroids exposure with onset of puberty in Chinese girls. *Environ Pollut* 2017b;**227**:606–612.
- Zhang Y, Cao Y, Shi H, Jiang X, Zhao Y, Fang X, Xie C. Could exposure to phthalates speed up or delay pubertal onset and development? A 1.5-year follow-up of a school-based population. *Environ Int* 2015;**83**:41–49.